

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 311/16, 493/10, 311/82, 265/38, C12Q 1/26		A1	(11) International Publication Number: WO 00/35900
		(43) International Publication Date: 22 June 2000 (22.06.00)	
(21) International Application Number: PCT/US99/29439		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU; ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 10 December 1999 (10.12.99)			
(30) Priority Data: 60/112,252 14 December 1998 (14.12.98) US 09/301,395 28 April 1999 (28.04.99) US 09/301,525 28 April 1999 (28.04.99) US			
(71) Applicant: AURORA BIOSCIENCES CORPORATION [US/US]; 11010 Torreyana Road, San Diego, CA 92121 (US).			
(72) Inventors: MAKINGS, Lewis ; 337 Cerro Street, Encinitas, CA 92024 (US). ZLOKARNIK, Gregor ; 2754 Palomino Circle, La Jolla, CA 92037 (US).			
(74) Agent: HUNT, Dale, C. ; Knobbe, Martens, Olson & Bear, LLP, 16th floor, 620 Newport Center Drive, Newport Beach, CA 92660 (US).		<p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>	

(54) Title: OPTICAL MOLECULAR SENSORS FOR CYTOCHROME P450 ACTIVITY



(57) Abstract

The invention provides a compound, useful as an optical probe or sensor of the activity of at least one cytochrome P450 enzyme, and methods of using the compound to screen candidate drugs, and candidate drugs identified by these methods. The optical probe of the invention is a compound having the generic structure Y-L-Q, wherein Y is selected from the group consisting of Q as herein defined, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl; L is selected from the group of (-OCR²H)_p, wherein for each p, all R² are separately selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and p is a positive integer no greater than twelve; and Q is a chemical moiety that gives rise to optical properties in its hydroxy or hydroxylate, phenol or phenoxide form that are different from the optical properties that arise from its ether form. Most preferably, p is one, R² is hydrogen, and Q is the ether form of a phenoxide fluorophore.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

OPTICAL MOLECULAR SENSORS FOR CYTOCHROME P450 ACTIVITY

Statement of Government Rights

5 This invention was developed in part, under grant number 1R43GM60114-01 from the National Institutes of General Medicine Sciences. The government may have certain rights in this invention.

Background of the InventionField of the Invention

10 This invention relates to novel chemical compounds, useful as optical indicators of cytochrome P450 activity, and especially to fluorogenic indicators of cytochrome P450 activity. More specifically, the invention relates to ether-containing compounds of the generic structure Y-L-Q, and to methods for assaying substrates and inhibitors of cytochrome P450 enzymes using these compounds in traditional assay formats, as well as
15 in high and ultra high throughput screening formats.

Description of the Related Art

20 The cytochrome P450 enzyme (CYP450) family comprises oxidase enzymes involved in the xenobiotic metabolism of hydrophobic drugs, carcinogens, and other potentially toxic compounds and metabolites circulating in blood. It is known that the liver is the major organ for xenobiotic metabolism, containing high levels of the most important CYP450 mixed-function oxygenases. There are numerous human P450 enzyme sub-families, often termed "isozymes" or "isoforms." Those of the CYP 3A4, CYP 2D6, CYP 2C, CYP 2A1 and CYP 2E1 subfamilies are known to be important in drug metabolism.
25 See, e.g., Murray, M., 23 Clin. Pharmacokinetics 132-46 (1992). Of these isoforms, CYP 3A4 is by far the major isoform in liver and the small intestines, comprising 30% and 70% respectively of the total CYP450 protein in those tissues. Based primarily on *in vitro* studies, it has been estimated that the metabolism of 40% to 50% of all drugs used in humans involve CYP 3A4 catalyzed oxidations. See Thummel, K.E. & Wilkinson, G.R., *In Vitro and In Vivo Drug Interactions Involving Human CYP 3A*, 38 Ann. Rev. Pharmacol.
30 Toxicol., 389-430 (1998).

Efficient metabolism of a candidate drug by a CYP450 enzyme may lead to poor pharmacokinetic properties, while drug candidates that act as potent inhibitors of a CYP450 enzyme can cause undesirable drug-drug interactions when administered with another drug that interacts with the same CYP450. See, e.g., Peck, C. C. *et al.*, *Understanding Consequences of Concurrent Therapies*, 269 JAMA 1550-52 (1993). Accordingly, early, reliable indication that a candidate drug interacts with (*i.e.*, is a substrate or inhibitor of) a CYP450 may greatly shorten the discovery cycle of pharmaceutical research and development, and thus may reduce the time required to market the candidate drug. Consequently, such earlier-available, reliable CYP450 pharmacokinetic information may result in greatly reduced drug development costs and/or increased profits from earlier market entrance. Furthermore, such earlier-available, reliable CYP450 pharmacokinetic information may allow a candidate drug to reach the public sooner, at lower costs than otherwise feasible. Accordingly, extensive pharmacokinetic studies of drug interactions in humans have recently become an integral part of the pharmaceutical drug development and safety assessment process. See, e.g., Parkinson, A., 24 *Toxicological Pathology* 45-57 (1996). Methodologies are therefore desired that will allow for (1) the more rapid acquisition of information about drug candidate interactions with CYP450 enzymes, earlier in the drug discovery process than presently feasible, and hence will allow for (2) the earlier elimination of unsuitable compounds and chemical series from further development efforts.

The need for information regarding drug candidate/CYP450 interactions has created a concurrent need for assays sensitive enough to test, in a cost-effective manner, vast arrays of compounds for interactions with the major human CYP450 enzymes involved in drug metabolism. Certain known techniques, including (1) CYP450 inhibition assays in which the metabolism of known CYP450 metabolite in the presence of the test compound, followed by quenching of the enzyme reaction and analysis of the extent of metabolism, (2) CYP450 metabolism of radioactively labeled test compound analogues, and (3) *in vivo* "cassette" dosing of animals (usually rats, dogs, or monkeys), see Berman, J. *et al.*, *Simultaneous Pharmacokinetic Screening of a Mixture of Compounds in the Dog using API LC/MS/MS Analysis for Increased Throughput*, 40 *J. Medicinal Chemistry*, 827-29 (1997), are not amenable to adaptation to miniaturization, or to the other requirements of high or ultra high throughput screening.

However, optical assays employing, for example, chromophores or luminescent phenols, and especially fluorescence-based assays are amenable to adaptation to miniaturization and high or ultra high throughput screening. Particularly, fluorescence-based assays have been used in pharmacokinetic studies of drug interactions in humans, more particularly in assays involving human hepatocyte cultures, where the number of available cells is severely limited. See Donato, M. T. *et al.*, 213 Anal. Biochem. 29-33 (1993).

Specifically, fluorogenic cytochrome P450 substrates have been commercially available for a number of years from, for example, Molecular Probes, Inc. (Eugene Oregon), SIGMA (St. Louis, Missouri), and more recently, GENTEST Corp. (Woburn, Massachusetts). Generally, these known fluorogenic CYP450 substrates are ether derivatives of well-known phenoxide type fluorophores, including: 7-hydroxycoumarin, fluorescein, and resorufin. Thus, generally, the CYP450 enzymes will catalyze a dealkylation reaction and convert the relatively non-fluorescent ether substrate into a relatively more highly-fluorescent phenoxide-containing product.

However, even the most recently developed fluorogenic CYP450 substrates either have relatively poor kinetics, or the enzymatic products do not have the desired physical and optical properties to allow reduction of the amount of enzyme needed to levels that would make large scale screening affordable and feasible. More specifically, these fluorogenic CYP450 substrates exhibit relatively poor turnover rates, poor aqueous solubility, low extinction coefficients and quantum yields, and/or weak fluorescence of the resultant phenolic dye. Furthermore, certain of these fluorogenic CYP450 substrates are excited in the ultraviolet, as opposed to visible, spectrum and therefore their signals are often masked by background stemming from the unreacted test compound. Finally, most of these fluorogenic CYP450 substrates are not specific for the CYP450 isozyme they are meant to detect, and therefore cannot be used for measurement in human liver microsomal preparations, a preferred analytical method that avoids potential artifacts caused by the alternative method of using an insect cell microsomal preparation. See Palamanda J.R. *et al.*, *Validation of a rapid microtiter plate assay to conduct cytochrome P450 2D6 enzyme inhibition studies*, 3 Drug Discovery Today 466-470 (1998). For these and other reasons, there exists an unfulfilled need for optical, and especially fluorogenic, CYP450 substrates that exhibit CYP450 isozyme-specificity, improved kinetics, and yield enzymatic products

having improved physical and optical properties for use in the screening of CYP450/drug candidate interactions, especially for use in high or ultra high throughput screening, and as part of the drug discovery process.

5

Summary of the Invention

The invention provides a compound, useful as an optical probe, modulator or sensor of the activity of at least one cytochrome P450 enzyme. The optical probe of the invention is a compound having the generic structure Y-L-Q, wherein Y is selected from the group consisting of Q as herein defined (such that the probe has the general structure Q-L'-Q), and
10 saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl; L is selected from the group of (-OCR²H)_p-, (-O(substituted ortho-phenyl)CR²H)_p-,
15 (-O(substituted meta-phenyl)CR²H)_p-, and (-O(substituted para-phenyl)CR²H)_p-, and L' is selected from the group of -(CR⁴H)(-OCR²H)_p-, -(CR⁴H)(-O(substituted ortho-phenyl)CR²H)_p-, -(CR⁴H)(-O(substituted meta-phenyl)CR²H)_p-, and -(CR⁴H)(-O(substituted para-phenyl)CR²H)_p-, wherein for each p, each R² is separately selected from the group consisting of a hydrogen atom, saturated C₁-
20 C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl and p is a positive integer no greater than twelve; and Q is a chemical moiety that gives rise
25 to optical properties in its hydroxy or hydroxylate, phenol or phenoxide form that are different from the optical properties that arise from its ether form. Most preferably, p is one, R² is hydrogen, and Q is the ether form of a phenoxide fluorophore. Most preferably, the ether form of the phenoxide fluorophore are ether derivatives of the well-known phenoxide type fluorophores 7-hydroxycoumarin, fluorescein, and resorufin.

30

The invention also provides methods for using the optical sensor compounds of the invention to determine whether a candidate drug, or class of candidate drugs, is a CYP450 substrate and/or whether the candidate drug, or class of candidate drugs, is a CYP450 inhibitor, and related methods for selecting a candidate drug, and for formulating and

administering that drug, having determined that the drug will not be metabolized by at least one CYP450 enzyme and/or that the drug will not act as an inhibitor of at least one CYP450 enzyme, and, thus, having determined that the drug will not, respectively, be too efficiently metabolized by a CYP 450 enzyme and/or elicit an unfavorable drug-drug interaction. Methods of selecting the candidate drug of the present invention may be by conventional methods or may be part of high or ultra high throughput screening of libraries of drug candidates.

Brief Description of the Drawings

The accompanying drawings, which are incorporated in and form part of the specification, merely illustrate embodiments of the present invention. Together with the remainder of the specification, they are meant to serve to explain the principles of the invention to those of skill in the art. In the drawings:

FIG. 1 illustrates Reaction Scheme 1, which shows a reaction mechanism for the CYP450 dealkylation of a currently-available fluorogenic CYP450 substrate, phenoxazone.

FIG. 2 illustrates Reaction Scheme 2, which shows a generic structure of the optical CYP450 substrate/sensor of the present invention, and the CYP450-catalyzed hydroxylation reaction.

FIG. 3 illustrates Reaction Scheme 3, which compares the hydroxylation reaction that may lead to a free phenolic dye of a known optical CYP450 sensor (top) and an optical CYP450 sensor compound of the present invention (bottom).

FIG. 4 illustrates a plot of the rate of resorufin ether conversion by CYP 3A4 as a function of CYP450 substrate/sensor concentration for the CYP450 substrate/sensor compounds of the invention, benzyloxymethylresorufin (BOMR) (circles) and n-octyloxymethylresorufin (OOMR) (diamonds), and as a function of resorufin benzyl ether (BR) (triangles).

FIG. 5 illustrates a plot of percent CYP 3A4 inhibition as a function of the presence of selected inhibitors and drug substrates of CYP 3A4, and demonstrates the effect that these inhibitors (cross-hatched bars) and drug substrates (diagonal striped bars) had on the turnover rate of a compound of the invention, benzyloxymethyl ether (BOMR), by the CYP 3A4 enzyme. This figure illustrates that inhibitors depress the rate of BOMR turnover

more than approximately 50%, while drug substrates slow the rate of BOMR turnover up to approximately 30%.

FIG. 6 illustrates a plot of percent CYP 2C19 inhibition as a function of the presence of various drugs at 10 μ M concentrations that interact with CYP 2C19, and demonstrates that 7-benzyloxymethoxy-3-cyanocoumarin (BOMCC) may be used as an optical CYP450 sensor to detect candidate drugs that interact with CYP 2C19.

FIG. 7 illustrates a plot of percent CYP 2C9 inhibition as a function of the presence of various drugs at 10 μ M concentrations that interact with CYP 2C9, and demonstrates that 7-benzyloxymethoxy-3-cyanocoumarin (BOMCC; dark bars) and octyloxymethyl-resorufin (OOMR; light bars) may be used as an optical CYP450 sensor to detect drugs that interact with CYP 2C9.

FIG. 8, 9, 10, 11 and 12 illustrate the improved signal over background of oxymethyl and oxyphenylmethyl linker containing sensors of this invention (solid traces) over prior commercially available substrates (broken lines). The cumulative background signal resulting from the addition of NADP+ and from addition of the substrate manifest itself by a fluorescence signal greater than one, 3 minutes following the substrate addition (second arrow indicates time of substrate addition). The later increase in signal, 4 minutes after 2nd addition and at later time points, is due to the enzymatic conversion of the substrate to the product. In Fig. 8, 9, 10, 11 and 12., enzymatic signal over reagent addition background, a measure of the performance of the assay, is greater for the oxymethyl-linker containing sensors of this invention (solid traces) than for prior commercially available substrates (broken lines). Figure 8. illustrates the superior signal to background of BOMR (solid trace) versus benzylresorufin (broken trace) with the CYP3A4 isozyme. Figure 9. illustrates the superior signal to background of BOMCC (solid trace) versus 7-Benzyloxy-4-trifluoromethylcoumarin (BFC, broken trace) with the CYP3A4 isozyme. Figure 10. illustrates the superior signal to background of MOBFC (solid trace) versus AMMC (Gentest) (broken trace) with the CYP2D6 isozyme. Figure 11. illustrates the superior signal to background of both OOMR (solid trace) and BOMCC (solid trace) versus 7-Methoxy-4-trifluoromethylcoumarin (MFC, broken trace) with the CYP2C9 isozyme. Figure 12. illustrates the superior signal to background of EOMCC (solid trace) versus 3-cyano-7-ethoxycoumarin (CEC, broken trace) with the CYP2C19 isozyme.

FIG. 13 illustrates the utility of oxymethyl-linker containing sensors in screening for CYP450 inhibitors. A random sample of 160 compounds purchased from Chembridge was screened for inhibition of CYP3A4 using BOMR as sensor to assess the degree of inhibition.

5 FIG. 14 illustrates the utility of oxymethyl-linker containing sensors in identifying structural motifs in chemicals that are associated with CYP3A4 inhibition.

FIG. 15 illustrates the utility of oxyphenylmethyl-linker containing sensors in screening for CYP450 inhibitors. A random sample of 240 compounds purchased from Chembridge was screened for inhibition of CYP2D6 using MOBFC as sensor to assess the
10 degree of inhibition.

FIG. 16 illustrates the utility of oxyphenylmethyl-linker containing sensors in identifying structural motifs in chemicals that are associated with CYP2D6 inhibition.


Detailed Description of the Preferred Embodiment

15 Members of the cytochrome P450 enzyme (CYP) family primarily catalyze epoxidation and hydroxylation reactions. Hydroxylation of a fluorogenic phenoxide ether liberates the free phenoxide which is readily detected by virtue of its fluorescence. The mechanisms of CYP450-catalyzed dealkylation reactions have been extensively studied and can be envisioned to proceed via the route depicted in Reaction Scheme 1, as illustrated in
20 FIG. 1. See Groves, J.T. et al., *Models and Mechanisms of Cytochrome P450 Action*, in "Cytochrome P450: Structure, Mechanism, Biochemistry," Plenum Press, 3-48, 1997. Experimental evidence suggests that the rate-limiting step in the reaction is the hydrogen abstraction reaction illustrated in the first step of Reaction Scheme 1, as illustrated in FIG. 1. Accordingly, a fluorogenic substrate with a faster turnover rate, especially with regard to
25 the rate-limiting step, may be desired to achieve the needs inherent in the art. A class of such substrates/sensors, the optical sensor compounds of the present invention, is provided, wherein the abstraction of any of the additional hydrogen atoms still generates a free compound in its hydroxy or hydroxylate, usually phenoxide, form which exhibits superior optical properties than the compound in its ether form.

30 The present invention provides, in a preferred embodiment, for the "insertion" of an oxymethyl linker between the fluorophore and the reactive ether moiety attached to the leaving group. Such an "insertion" is accomplished, according to, for example, the

synthetic methods of EXAMPLES 1 through 7, which are preferred methods of preparing the optical CYP450 sensors of the present invention.

Generally, as will be appreciated upon review of EXAMPLES 1 through 7 by persons of skill in the art, compounds of the present invention may be synthesized according to the following reaction scheme:



wherein X is a suitable leaving group, for example a halogen atom, a tosyl group, a mesyl group, a triflate group, and wherein the reaction is carried out in the presence of, preferably, DMF/K₂CO₃, diisopropylethylamine/DMF at temperatures at or slightly above the freezing point of water; wherein Q is a compound which exhibits superior optical properties in its hydroxy or hydroxylate, typically but not exclusively phenoxide, form than it does as in its ether form, and is preferably a fluorophore or a chromophore, and is most preferably a fluorophore selected from the group consisting of 7-hydroxycoumarin, resorufin, and the known phenoxide fluorophores; and R³ is selected from the group consisting of Q as herein defined, R¹ of the known fluorogenic cytochrome P450 substrates, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl.

Furthermore, and more generally descriptive of compounds of the present invention, one of the methyl protons of the linker may be replaced by a distinct chemical group, R², wherein R² is selected from the group consisting of saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl groups. Furthermore, multiple, linked oxymethyl, or more generally, multiple OCR²H, groups may form the linker of the CYP450 sensor of the invention. In a multimeric linker the R² groups are selected independently from each other. For example, a linker denoted (CR²H)_p with p=3 has following structure: -(OCR²H)-(OCR²H)-(OCR²H)-, in which R² and R² and R² are independently selected from the group consisting of a hydrogen atom, saturated C₁-

C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl.

The present invention therefore provides compounds, useful as optical probes for quantifying the activity of at least one cytochrome P450 enzyme; said compound having the generic structure Y-L-Q wherein:

Y is selected from the group consisting of (i) Q as herein defined, so long as L is L' as herein defined, and (ii) the group consisting of saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl; substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl.

L is selected from the group of (-OCR²H)_p-, (-O(substituted ortho-phenyl)CR²H)_p-, (-O(substituted meta-phenyl)CR²H)_p-, and (-O(substituted para-phenyl)CR²H)_p-, wherein for each p, each R² is separately selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, and p is a positive integer no greater than twelve. When Y is selected from Q as herein defined, L is L', wherein L' is selected from the group of -(CR⁴H)(-OCR²H)_p-, -(CR⁴H)(-O(substituted ortho-phenyl)CR²H)_p-, -(CR⁴H)(-O(substituted meta-phenyl)CR²H)_p-, and -(CR⁴H)(-O(substituted para-phenyl)CR²H)_p-, wherein each R² and R⁴ is separately selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, and p is a positive integer no greater than twelve.

Use of the structures Q-L'-Q of the CYP450 sensor of the invention has the advantage of yielding two, instead of one, optical Q moieties in the hydroxy or hydroxylate form upon interaction of the optical sensor of the invention with at least one CYP450 enzyme.

5 The terms substituted ortho-phenyl, substituted meta-phenyl, and substituted para-phenyl refer to a phenyl that is part of the linker connecting Y with Q in which ortho, meta, and para refer to positions of the carbons in the phenyl ring that serve as the attachment for Y and Q. Ortho substituted refers to attachment of Y and Q via adjacent carbons in the phenyl ring, meta substituted refers to attachment of Y and Q by carbons spaced by one
10 carbon on the phenyl ring, and para substitution refers to the attachment of Y and Q on the phenyl ring by carbons that are spaced by two carbons on the phenyl ring. When used in defining additional substitution of the phenyl ring in the oxyphenylmethyl linker the term substituted refers to the substitution of the remaining carbons not involved in attachment of Y and Q on the phenyl ring


15 The term "substituted" means any substitution of a hydrogen atom with a functional group. Functional groups are selected from the group consisting of a halogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, -SR_S, -OR_O, -NR_{n1}R_{n2}, -N⁺R_{n1}R_{n2}R_{n3}, -N=N-R_{n1}, -P⁺R_{n1}R_{n2}R_{n3}, -COR_C, -C(=NOR_O)R_C, -CSR_C, -OCOR_C, -
20 OCONR_{n1}R_{n2}, -OCO₂R_C, -CONR_{n1}R_{n2}, -C(=N)NR_{n1}R_{n2}, -CO₂R_O, -SO₂NR_{n1}R_{n2}, -SO₃R_O, -SO₂R_O, -PO(OR_O)₂, -NR_{n1}CSNR_{n2}R_{n3}, -NR_{n1}C(=N)NR_{n2}R_{n3}, -NR_{n1}CONR_{n2}R_{n3}, -NR_{n1}COR_C and -NR_{n1}S(=O)₂R_S. Substituents R_{n1}, R_{n2}, R_{n3}, R_O and R_S are each separately selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and
25 may constitute parts of an aliphatic or aromatic heterocycle. R_C is selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and cyano. Also, when used in the context of defining Y and R², the term "substituted" means any substitution of a hydrogen with a functional group as defined
30 herein so long as hetero-atom substitution does not occur at the α-carbon.

The term "quencher" refers to a chromophoric molecule or part of a compound, which is capable of reducing the emission from a fluorescent donor when attached to the


donor. Quenching may occur by any of several mechanisms including fluorescence resonance energy transfer, photoinduced electron transfer, paramagnetic enhancement of intersystem crossing, Dexter exchange coupling, and exciton coupling such as the formation of dark complexes.

5 The term "acceptor" refers to a quencher that operates via energy transfer. Acceptors may re-emit the transferred energy as fluorescence and are "acceptor fluorescent moieties". Examples of acceptors include coumarins and related fluorophores, xanthenes such as fluoresceins, rhodols, and rhodamines, resorufins, cyanines, difluoroboradiazaindacenes, and phthalocyanines. Other chemical classes of acceptors
10 generally do not re-emit the transferred energy as light. Examples include indigos, benzoquinones, anthraquinones, azo compounds, nitro compounds, indoanilines, and di- and triphenylmethanes.


Q is attached to L through an ether linkage via the oxygen indicated by the arrow, and has a structure selected from the group consisting of the following structures:




I




II




III




IV




V




VI




VII




VIII




IX




X




XI



XII




XIII




XIV


5




XV




XVI




XVII



XVIII




XIX




XX


5




XXI




XXII




XXIII



XXIV




XXV




XXVI

10



XXVII'



XXVIII

wherein:

m is a positive integer no greater than five;

R_a , R_b , R_c , R_d , R_e , R_f , R_g , R_h , R_i , R_j , R_k , and R_l are each separately selected from the group consisting of a hydrogen atom, a halogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, azido, $-SR_s$, $-OR_o$, $-NR_{n1}R_{n2}$, $-N^+R_{n1}R_{n2}R_{n3}$, $-N=N-R_{n1}$, $-P^+R_{n1}R_{n2}R_{n3}$, $-COR_c$, $-C(=NOR_o)R_c$, $-CSR_c$, $-OCOR_c$, $-OCONR_{n1}R_{n2}$, $-OCO_2R_c$, $-CONR_{n1}R_{n2}$, $-C(=N)NR_{n1}R_{n2}$, $-CO_2R_o$, $-SO_2NR_{n1}R_{n2}$, $-SO_3R_o$, $-SO_2R_o$, $-PO(OR_o)_2$, $-NR_{n1}CSNR_{n2}R_{n3}$, $-NR_{n1}C(=N)NR_{n2}R_{n3}$, $-NR_{n1}CONR_{n2}R_{n3}$, $-NR_{n1}COR_c$ and $-NR_{n1}S(=O)_2R_s$;

R_{n1} , R_{n2} , R_{n3} , R_o and R_s are each separately selected from the group consisting of a hydrogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of an aliphatic or aromatic heterocycle;

R_c is selected from the group consisting of a hydrogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano and may constitute parts of an aliphatic or aromatic homo- or heterocycle;

A is selected from the group consisting of an oxygen atom, a sulfur atom, SO , SO_2 , $C(CH_3)_2$ and $C(CF_3)_2$;

E and E' are separately selected from the group consisting of an oxygen atom, a sulfur atom and NR_{n1} ;

G is selected from the group consisting of an oxygen atom, a sulfur atom, and $NR_{n1}R_{n2}$, wherein if G is selected from $NR_{n1}R_{n2}$, G and R_c , as well as G and R_d , may constitute parts of a heterocycle; and

T is selected from the group consisting of an oxygen atom and NR_{n1} .

As will be appreciated by reference to the presently presented examples of the optical CYP450 sensors of the present invention, the preferred optical sensors of the present invention are fluorogenic CYP450 sensors wherein Q is the ether form of a phenoxide fluorophore. Furthermore, the forgoing examples of Q are meant to highlight the point that Q may be any chemical structure, so long as Q is a chemical means for generating an altered optical signal via cleavage of a C-O bond. Those of skill in the art will recognize variations to the structures of Q herein described to achieve this function. Furthermore, the

function of generating an altered optical signal via cleavage of a C-O bond may be achieved by releasing a dye upon cleavage of a C-O bond or, more preferably, by releasing a fluorescent dye upon cleavage of a C-O bond, and even more preferably, by releasing a phenolic fluorescent dye upon cleavage of a C-O bond. Preferably, the altered optical signal is an enhanced optical signal.

In the case where Q is an ether form of a fluorophore, Y may act as a quencher. In this case, CYP450 activity is detected by an increase in fluorescence from Q, which is due to the loss of quenching of its fluorescence by Y. If fluorescence quenching by Y occurs via fluorescence resonance energy transfer, then Y is referred to as an acceptor. In the case where Q is an ether form of a fluorophore and Y acts as a quencher, attachment of Y-L to Q can be by substitution of any hydrogen on the fluorophore by Y-L-O-, the O denoting an oxygen atom. In this case Q can be any fluorophore, the ether form of which being formed by substitution of one or more fluorophore hydrogen atoms by Y-L-O-. See U.S. Patent No.: 5,741,657 to Tsien and Zlokarnik (issued April 21, 1998), which is incorporated by reference herein.

In the optical CYP450 sensors of the present invention, Y is preferably selected from C₁-C₈ alkyl, C₁-C₈ alkenyl, substituted C₂-C₈ alkyl, substituted C₂-C₈ alkenyl, alkoxyalkyl, aryl, substituted aryl, tertiary and quarternary aminoalkyl and guanidinium groups. Among the aryls and substituted aryls, benzyl, and substituted benzyl groups are most preferred. Most preferably, Y is selected from methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, octyl and benzyl.

Preferably, L is selected from the group consisting of -(OCR²H)_p- and -(O-para-phenyl-CR²H)_p- wherein R² is a hydrogen atom or methyl, and p equals either one or two. Most preferably R² is a hydrogen atom and p equals one.

Preferably, Q is a fluorophore. More preferably, Q is selected from the group consisting of 7-hydroxycoumarin, resorufin, fluorescein and other phenoxide fluorophores. Nevertheless Q may be a chromophore, so long as it exhibits optical properties in its hydroxy or hydroxylate form, *e.g.*, after interaction with an active CYP450 enzyme, that differ from its ether form, *e.g.*, in its unreacted state. Most generally, Q is a chemical moiety that exhibits optical properties in its free hydroxy or its hydroxylate, usually phenoxide, form that are different from the optical properties that it exhibits in its ether

form. Suitable structures of Q, as used herein, may also be found in U.S. Patent No. 5,741,657, which is incorporated by reference herein.

The optical CYP450 sensor compounds of the present invention may be used to determine CYP450 activities by a variety of optical signals, including for example, in the context of (a) the CYP450-catalyzed formation of chromogenic or fluorogenic or luminescent phenols, (b) the CYP450-catalyzed formation of chromogenic or fluorogenic precipitates, (c) the CYP450-catalyzed light generation from conversion of a phenolic dioxetane substrate, (d) the CYP450-catalyzed liberation of a salicylate or other phenolic ligand detectable by heavy metal chelate formation to give a colored, fluorescent, phosphorescent or electrochemiluminescent product, and (e) the CYP450-catalyzed liberation of a sensitizer for light generation by peroxide/luminol, and (f) the CYP450-catalyzed liberation of a substrate suitable for secondary enzyme detection (e.g., the liberation of a luciferin, which may be detected by a luciferase).

As used herein, the terms "halogen" and "halogen atom" refer to any one of the radio-stable atoms of column 17 of the Periodic Table of the Elements, i.e., fluorine, chlorine, bromine, or iodine, with fluorine and chlorine being most preferred.

As used herein, the term "alkyl" means any unbranched, branched or cyclic, saturated hydrocarbon, with C₁-C₈ unbranched, saturated, unsubstituted hydrocarbons being preferred, and with methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, n-pentyl and n-octyl being most preferred.

The term "substituted alkyl" means any unbranched, branched or cyclic, substituted saturated hydrocarbon substituted with one or more functional groups. Functional groups are selected from the group consisting of a halogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, -SR_S, -OR_O, -NR_{n1}R_{n2}, -N⁺R_{n1}R_{n2}R_{n3}, -P⁺R_{n1}R_{n2}R_{n3}, -COR_C, -C(=NOR_O)R_C, -CSR_C, -OCOR_C, -OCONR_{n1}R_{n2}, -OCO₂R_C, -CONR_{n1}R_{n2}, -C(=N)NR_{n1}R_{n2}, -CO₂R_O, -SO₂NR_{n1}R_{n2}, -SO₃R_O, -SO₂R_O, -PO(OR_O)₂, -NR_{n1}CSNR_{n2}R_{n3}, -NR_{n1}C(=N)NR_{n2}R_{n3}, -NR_{n1}CONR_{n2}R_{n3}, -NR_{n1}COR_C and -NR_{n1}S(=O)₂R_S. Substituents R_{n1}, R_{n2}, R_{n3}, R_O and R_S are each separately selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of an aliphatic or aromatic heterocycle. R_C is selected from the group consisting of a

hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and cyano. When used in the context of defining Y and R², the term "substituted alkyl" means any unbranched or branched, substituted saturated hydrocarbon, so long as hetero-atom substitution does not occur at the α-carbon.

The term "alkenyl" means any unbranched, branched or cyclic, substituted or unsubstituted, unsaturated hydrocarbon, with C₁-C₈ unbranched, mono-unsaturated and di-unsaturated being preferred. The term "substituted alkenyl" means any unbranched or branched, substituted unsaturated hydrocarbon substituted with one or more functional groups. Functional groups are selected from the group consisting of a halogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, , -SR_S, -OR_O, -NR_{n1}R_{n2}, -N⁺R_{n1}R_{n2}R_{n3}, -P⁺R_{n1}R_{n2}R_{n3}, -COR_C, -C(=NOR_O)R_C, -CSR_C, -OCOR_C, -OCONR_{n1}R_{n2}, -OCO₂R_C, -CONR_{n1}R_{n2}, -C(=N)NR_{n1}R_{n2}, -CO₂R_O, -SO₂NR_{n1}R_{n2}, -SO₃R_O, -SO₂R_O, -PO(OR_O)₂, -NR_{n1}CSNR_{n2}R_{n3}, -NR_{n1}C(=N)NR_{n2}R_{n3}, -NR_{n1}CONR_{n2}R_{n3}, -NR_{n1}COR_C and -NR_{n1}S(=O)₂R_S. Substituents R_{n1}, R_{n2}, R_{n3}, R_O and R_S are each separately selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of an aliphatic or aromatic heterocycle. R_C is selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano and may constitute parts of an aliphatic or aromatic homo- or heterocycle. When used in the context of defining Y and R², the term "substituted alkenyl" means any unbranched or branched, substituted unsaturated hydrocarbon, so long as neither the carbon-carbon double bond, nor heteroatom substitution occurs at the α-carbon.

The terms "aryl," "substituted aryl," "heteroaryl," and "substituted heteroaryl" refer to aromatic hydrocarbon rings, preferably having five or six atoms comprising the ring. The term "substituted aryl" includes mono and poly-substituted aryls, substituted with functional groups selected from the group of a halogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, azido, -SR_S, -OR_O, -NR_{n1}R_{n2}, -N⁺R_{n1}R_{n2}R_{n3}, -N=N-R_{n1}, -P⁺R_{n1}R_{n2}R_{n3}, -COR_C, -C(=NOR_O)R_C, -CSR_C, -OCOR_C,

-OCONR_{n1}R_{n2}, -OCO₂R_C, -CONR_{n1}R_{n2}, -C(=N)NR_{n1}R_{n2}, -CO₂R_O, -SO₂NR_{n1}R_{n2}, -SO₃R_O, -SO₂R_O, -PO(OR_O)₂, -NR_{n1}CSNR_{n2}R_{n3}, -NR_{n1}C(=N)NR_{n2}R_{n3}, -NR_{n1}CONR_{n2}R_{n3}, -NR_{n1}COR_C and -NR_{n1}S(=O)₂R_S. Substituents R_{n1}, R_{n2}, R_{n3}, R_O and R_S are each separately selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of an aliphatic or aromatic heterocycle. R_C is selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, and may constitute parts of an aliphatic or aromatic homo- or heterocycle. "Heteroaryl" and "substituted heteroaryl," refer to aromatic hydrocarbon rings in which at least one heteroatom, *e.g.*, oxygen, sulfur, or nitrogen atom, is in the ring along with at least one carbon atom.

The terms substituted ortho-phenyl, substituted meta-phenyl, and substituted para-phenyl refer to a phenyl that is part of the linker connecting Y with Q in which ortho, meta and para refer to position of the carbons in the phenyl ring that serve as the attachment for Y and Q. Ortho substituted refers to attachment of Y and Q via adjacent carbons in the phenyl ring, meta substituted refers to attachment of Y and Q by carbons spaced by one carbon on the phenyl ring and para substitution refers to the attachment of Y and Q on the phenyl ring by carbons that are spaced by two carbons on the phenyl ring. When used in defining additional substitution of the phenyl ring in the oxyphenylmethyl linker, the term "substituted" refers to the substitution of hydrogens on the remaining carbons not involved in attachment of Y and Q on the phenyl ring. The term "substituted phenyl" includes mono and poly-substituted phenyls, substituted with functional groups. Functional groups are selected from the group consisting of a halogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, azido, nitro, -, -SR_S, -OR_O, -NR_{n1}R_{n2}, -N⁺R_{n1}R_{n2}R_{n3}, -P⁺R_{n1}R_{n2}R_{n3}, -COR_C, -C(=NOR_O)R_C, -CSR_C, -OCOR_C, -OCONR_{n1}R_{n2}, -OCO₂R_C, -CONR_{n1}R_{n2}, -C(=N)NR_{n1}R_{n2}, -CO₂R_O, -SO₂NR_{n1}R_{n2}, -SO₃R_O, -SO₂R_O, -PO(OR_O)₂, -NR_{n1}CSNR_{n2}R_{n3}, -NR_{n1}C(=N)NR_{n2}R_{n3}, -NR_{n1}CONR_{n2}R_{n3}, -NR_{n1}COR_C and -NR_{n1}S(=O)₂R_S. Substituents R_{n1}, R_{n2}, R_{n3}, R_O and R_S are each separately selected from the group consisting of a hydrogen atom, C₁-C₂₀ alkyl, substituted C₁-C₂₀ alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of

an aliphatic or aromatic heterocycle. R_C is selected from the group consisting of a hydrogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, perhalogenated alkyl, cycloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, and may constitute parts of an aliphatic or aromatic homo- or heterocycle.

5 The terms "fluorogenic CYP450 substrate" generally refers to any compound that, upon interacting with a CYP450 enzyme, exhibits superior fluorescence properties than the compound exhibited prior to interacting with the CYP450 enzyme. As used herein, the terms "optical probe" and "optical sensor," are synonymous, each referring to a compound that can be used to assay an activity that catalyzes the conversion of the ether form of the
10 compound to the hydroxy or hydroxylate, usually phenoxide, form of the compound by virtue of the fact that each contains a chemical moiety that exhibits optical properties in its hydroxy or hydroxylate, usually its phenoxide, form, that are distinct from, and preferably superior to, the optical properties that the chemical moiety exhibits as an ether. The terms "optical CYP450 probe." and "optical CYP450 sensor" are synonymous; each is a broader
15 term than "fluorogenic CYP450 substrate"; each refers to a compound that may be used to assay the presence and, especially where a CYP450 inhibitor may be present, the activity of at least one CYP450 enzyme by virtue of the fact that each contains a chemical moiety that exhibits optical properties in its hydroxy or hydroxylate, usually its phenoxide, form, that are distinct from, and preferably superior to, the optical properties that the chemical moiety
20 exhibits as an ether. By virtue of the fact that at least one CYP450 enzyme will catalyze the conversion of the ether form to the hydroxy or hydroxylate, usually phenoxide, form, these optical probes or sensors may be used to assay the presence and activity of at least one CYP450 enzyme.


25 The term "reagent compound" refers to the compounds of the invention, as herein described, especially the compounds of general structure Y-L-Q. The term "candidate compound" is a term broader than the terms "candidate drug" and "candidate modulator," as those term are used herein, and refers to any compound, of whatever origin, suitable for being screened for its activity as a substrate or inhibitor of a CYP450 enzyme according to the methods of the present invention.

30 Within the present invention, long wavelength fluorescence dyes are preferred over dyes that are excited in the UV, but all fluorescence dyes, as well as dyes that are excited in

the UV or IR, are useful as Q in the optical sensor compounds and the methods of the invention.

Many screening compound libraries often contain fluorescent compounds. Typically the fluorescent compounds in libraries have absorbances in the UV or short wavelength visible portion of the spectrum. Thus, for many fluorescent assays, longer wavelength reporter molecules usually result in assays that have lower background and less interference. In addition, compounds of the present invention preferable have improved solubility in both water and acetonitrile compared to the most closely related CYP450 substrates currently available. Aqueous solubility is important, as 1-20 μ M substrate concentrations are needed to lead to a strong fluorescence signal in the assay. Good solubility in acetonitrile (1-10mM) allows the delivery of the hydrophobic substrate molecules into the aqueous assay medium in small volumes. Acetonitrile is a preferred solvent, as it does not inhibit CYP450 at concentrations up to 2%. Other solvents, such as DMSO and ethanol, typically used to deliver hydrophobic molecules into the aqueous assay medium do inhibit the activity of most CYP450 enzymes at lower concentrations and are therefore not preferred substrate delivery. However, as is known, CYP450 and related compounds do tolerate DMSO at concentrations up to 0.5%, permitting delivery of test compounds to the assay medium in this solvent.


With reference to the following structure, it has been demonstrated that the introduction of an oxymethyl spacer ($R^2 = H$) between the moiety R^1 and the phenolic dye of currently-available fluorogenic CYP450 sensors tends to increase the efficiency (k_{cat}/K_m) of turnover by many CYP450 enzymes and related compounds. R^1 of currently available fluorogenic CYP450 sensors are alkyl or substituted methyl, with the Substituent being an aryl group or a steroid, *see* U.S. Patent 5,110,725. An optical CYP450 probe of the present invention, shown in the following structure, illustrates this "insertion" to provide the optical CYP450 sensor compounds of the present invention.



This improved turnover, and improved optical properties, have been demonstrated for a variety of structurally-distinct substrates. Furthermore, solubilities of sensors of the invention in acetonitrile, as well as water, are excellent, overcoming one of the above-mentioned limitations of the currently available fluorogenic CYP450 substrates. Accordingly, the above structure illustrates optical CYP450 sensor compounds of the present invention, wherein R^1 is a structure as herein defined for Y. Thus, R^1 in the above structure of CYP450 sensor compounds of the present invention is selected from a group consisting of all Y as herein defined. However, the groups corresponding to R^1 that are found on presently, commercially-available phenol CYP450 ether substrates, are but a subset of Y as herein defined; compounds having the linker of the present invention and employing groups corresponding to R^1 that are found in presently, commercially-available phenol CYP450 ether substrates-compounds lacking the linker of the present invention-exhibit improved physical and optical properties with respect to presently, commercially-available phenol CYP450 ether substrates.

R^2 in the above structure of CYP450 sensor compounds of the present invention is selected from the group consisting of a hydrogen atom, saturated C_1 - C_{20} alkyl, unsaturated C_1 - C_{20} alkenyl, unsaturated C_1 - C_{20} alkynyl, substituted saturated C_1 - C_{20} alkyl, substituted unsaturated C_1 - C_{20} alkenyl, substituted unsaturated C_1 - C_{20} alkynyl, C_1 - C_{20} cycloalkyl, C_1 - C_{20} cycloalkenyl, substituted saturated C_1 - C_{20} cycloalkyl, substituted unsaturated C_1 - C_{20} cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl groups, as those terms are herein defined.

Furthermore, it has been demonstrated that the introduction of an oxyphenylmethyl spacer ($R^2 = H$) between the moiety R^1 and the phenolic dye of currently-available fluorogenic CYP450 sensors also tends to increase the efficiency (k_{cat}/K_m) of turnover by CYP450 enzymes and related compounds.



Accordingly, the above structure illustrates oxyphenylmethyl containing optical CYP450 sensor compounds of the present invention, wherein R¹ is a structure as herein defined for Y. Thus, R¹ in the above structure of CYP450 sensor compounds of the present invention is selected from a group consisting of all Y as herein defined. However, the groups
5 corresponding to R¹ that are found on presently, commercially-available phenol CYP450 ether substrates, are but a subset of Y as herein defined; compounds having the linker of the present invention and employing groups corresponding to R¹ that are found in presently, commercially-available phenol CYP450 ether substrates-compounds lacking the linker of the present invention-exhibit improved physical and optical properties with respect to
10 presently, commercially-available phenol CYP450 ether substrates.

R² in the above structure of CYP450 sensor compounds of the present invention is selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted
15 unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl, as those terms are herein defined.

According to present invention, FIG. 2 illustrates Reaction Scheme 2, which shows a generic structure of the optical CYP450 substrate/sensor of the present invention, and the
20 CYP450-catalyzed hydroxylation reaction. FIG. 3 illustrates Reaction Scheme 3, which compares the hydroxylation reaction that may lead to a free phenolic dye of a known optical CYP450 sensor (top) and an optical CYP450 sensor compound of the present invention (bottom).

As herein described, candidate drugs can be screened and evaluated for their
25 activities as substrates of or inhibitors of a CYP450 enzyme by using the optical CYP450 sensors of the present invention. A candidate drug may be determined to be an inhibitor or a substrate of a cytochrome P450 enzyme by contacting a cytochrome P450 enzyme with the candidate drug, under conditions suitable for interaction therebetween, providing at least one optical cytochrome P450 enzyme sensor, under conditions that would, in the
30 absence of an inhibitor or substrate of the cytochrome P450 enzyme, be suitable for interaction between the optical cytochrome P450 enzyme sensor and the cytochrome P450 enzyme, and detecting the presence of signal of a free phenolic dye, wherein the phenolic

dye would be, in the absence of an inhibitor of the cytochrome P450 enzyme, the product of the reaction between the cytochrome P450 enzyme and the optical cytochrome P450 enzyme sensor. Such efficient CYP450 substrates and inhibitors, as deemed appropriate by those of skill in the art, may be removed from a screening library where such efficient CYP450 substrates and inhibitors are not desired in the remainder of the screening for a candidate drug.

To distinguish between a substrate and an inhibitor of cytochrome P450 enzymes, typically, the candidate compound is incubated with at least one cytochrome P450 enzyme under conditions, which allow for metabolism of the candidate compound prior to providing the optical cytochrome P450 enzyme sensor under conditions that would, in the absence of an inhibitor or substrate of the cytochrome P450 enzyme, be suitable for interaction between the optical cytochrome P450 enzyme sensor and the cytochrome P450 enzyme. The resulting optical signal is compared to the one obtained from contacting a cytochrome P450 enzyme with the candidate drug, under conditions suitable for interaction therebetween, providing at least one optical cytochrome P450 enzyme sensor, under conditions that would, in the absence of an inhibitor of the cytochrome P450 enzyme, be suitable for interaction between the optical cytochrome P450 enzyme sensor and the cytochrome P450 enzyme. Metabolism of the candidate drug by a cytochrome P450 enzyme reduces its concentration in the assay medium and may lead to an apparent loss of cytochrome P450 inhibitory activity compared to conditions without metabolism of the compound which would indicate it was a substrate for the enzyme. An inhibitory compound that was not metabolized would show equal potency, irrespective of the time of addition of the optical cytochrome p450 enzyme substrate.

The following procedures may be used to then further screen, formulate, and administer the candidate drugs of the present invention. These drugs are within the present invention to the extent that they have not yet been identified as candidate drugs or modulators, and to the extent that they are identified as candidate drugs or modulators by means of using the optical sensors of the present invention.

In certain cases, a candidate drug may be determined to be a cytochrome P450 enzyme substrate of at least one cytochrome P450 enzyme, by selecting an optical cytochrome P450 enzyme sensor that is a derivative of the candidate drug; contacting a cytochrome P450 enzyme with the optical cytochrome P450 enzyme sensor under

conditions suitable for interaction therebetween, and detecting the absence of signal of a free phenolic dye, that would be the product of the reaction between the cytochrome P450 enzyme and the optical cytochrome P450 enzyme sensor.

Bioavailability and Toxicology of Candidate Modulators

5 Once identified, candidate drugs or modulators can be further evaluated for bioavailability and toxicological effects using known methods. *See* Lu, *Basic Toxicology, Fundamentals, Target Organs, and Risk Assessment*, Hemisphere Publishing Corp., Washington (1985); U.S. Patent Nos: 5,196,313 to Culbreth (issued March 23, 1993) and U.S. Patent No. 5,567,952 to Benet (issued October 22, 1996). For example, toxicology of
10 a candidate modulator can be established by determining *in vitro* toxicity towards a cell line, such as a mammalian *i.e.* human, cell line. Candidate modulators can be treated with, for example, tissue extracts, such as preparations of liver, such as microsomal preparations, to determine increased or decreased toxicological properties of the chemical after being metabolized by a whole organism. The results of these types of studies are often predictive
15 of toxicological properties of chemicals in animals, such as mammals, including humans.

Such bioavailability and toxicological methods can be performed as part of or as complimentary to the screening systems and methods of the present invention. Such methods include contacting a sample having a target with at least one photon producing agent, at least one photon reducing agent, and a test chemical. An optical signal from said
20 at least one photon producing agent is detected, wherein said optical signal is related to a toxicological activity. Bioavailability is any known in the art and can be detected, for example by measuring reporter genes that are activated during bioavailability criteria. Toxicological activity is any known in the art, such as apoptosis, cell lysis, crenation, cell death and the like. The toxicological activity can be measured using reporter genes that are
25 activated during toxicological activity or by cell lysis (*see* WO 98/13353, published 4/2/98). Preferred reporter genes produce a fluorescent or luminescent translational product (such as, for example, a Green Fluorescent Protein (*see*, for example, U.S. Patent No. 5,625,048 to Tsien et al., issued 4/29/98; U.S. Patent No. 5,777,079 to Tsien et al., issued 7/7/98; WO 96/23810 to Tsien, published 8/8/96; WO 97/28261, published 8/7/97; PCT/US97/12410, filed 7/16/97; PCT/US97/14595, filed 8/15/97)) or a translational
30 product that can produce a fluorescent or luminescent product (such as, for example, beta-lactamase, *e.g.*, U.S. Patent No. 5,741,657 to Tsien, issued 4/21/98, and WO 96/30540,

published 10/3/96), such as an enzymatic degradation product. Cell lysis can be detected in the present invention as a reduction in a fluorescence signal from at least one photon-producing agent within a cell in the presence of at least one photon reducing agent. Such toxicological determinations can be made using prokaryotic or eukaryotic cells, optionally using toxicological profiling, such as described in PCT/US94/00583, filed 1/21/94, German Patent No 69406772.5-08, issued 11/25/97; EPC 0680517, issued 11/12/94; U.S. Patent No. 5,589,337, issued 12/31/96; EPO 651825, issued 1/14/98; and U.S. Patent No. 5,585,232, issued 12/17/96).

Alternatively, or in addition to these *in vitro* studies, the bioavailability and toxicological properties of a candidate modulator in an animal model, such as mice, rats, rabbits, or monkeys, can be determined using established methods. See, Lu, *supra* (1985); and Creasey, *Drug Disposition in Humans, The Basis of Clinical Pharmacology*, Oxford University Press, Oxford (1979), Osweiler, *Toxicology*, Williams and Wilkins, Baltimore, MD (1995), Yang, *Toxicology of Chemical Mixtures, Case Studies, Mechanisms, and Novel Approaches*, Academic Press, Inc., San Diego, CA (1994), Burrell et al., *Toxicology of the Immune System: A Human Approach*, Van Nostrand Reinhold, Co. (1997), Niesink et al., *Toxicology: Principles and Applications*, CRC Press, Boca Raton, FL (1996). Depending on the toxicity, target organ, tissue, locus, and presumptive mechanism of the candidate modulator, the skilled artisan would not be burdened to determine appropriate doses, LD₅₀ values, routes of administration, and regimes that would be appropriate to determine the toxicological properties of the candidate modulator. In addition to animal models, human clinical trials can be performed following established procedures, such as those set forth by the United States Food and Drug Administration (USFDA) or equivalents of other governments. These toxicity studies provide the basis for determining the undesired effects of a candidate modulator *in vivo*.

Efficacy of Candidate Modulators

Efficacy of a candidate modulator can be established using several art recognized methods, such as *in vitro* methods, animal models, or human clinical trials, see, Creasey, *supra* (1979). Recognized *in vitro* models exist for several diseases or conditions. For example, the ability of a chemical to extend the life-span of HIV-infected cells *in vitro* is recognized as an acceptable model to identify chemicals expected to be efficacious to treat HIV infection or AIDS, see, Daluge et al., *Antimicro. Agents Chemother.* 41:1082-1093

(1995). Furthermore, the ability of cyclosporin A (CsA) to prevent proliferation of T-cells *in vitro* has been established as an acceptable model to identify chemicals expected to be efficacious as immunosuppressants, *see*, Suthanthiran et al., *supra*, (1996). For nearly every class of therapeutic, disease, or condition, an acceptable *in vitro* or animal model is available. Such models exist, for example, for gastro-intestinal disorders, cancers, cardiology, neurobiology, and immunology. In addition, these *in vitro* methods can use tissue extracts, such as preparations of liver, such as microsomal preparations, to provide a reliable indication of the effects of metabolism on the candidate modulator. Similarly, acceptable animal models may be used to establish efficacy of chemicals to treat various diseases or conditions. For example, the rabbit knee is an accepted model for testing chemicals for efficacy in treating arthritis. *See* Shaw and Lacy, *J. Bone Joint Surg. (Br)* 55:197-205 (1973)). Hydrocortisone, which is approved for use in humans to treat arthritis, is efficacious in this model which confirms the validity of this model. *See*, McDonough, *Phys. Ther.* 62:835-839 (1982). When choosing an appropriate model to determine efficacy of a candidate modulator, the skilled artisan can be guided by the state of the art to choose an appropriate model, dose, and route of administration, regime, and endpoint and as such would not be unduly burdened.

In addition to animal models, human clinical trials can be used to determine the efficacy of a candidate modulator in humans. The USFDA, or equivalent governmental agencies, have established procedures for such studies, *see, e.g.*, <http://www/fda.gov>.

Selectivity of Candidate Modulators

The *in vitro* and *in vivo* methods described above as part of the present invention also establish the selectivity of a candidate drug or modulator. It is recognized that chemicals can modulate a wide variety of biological processes or be selective. Panels of cells based on the present invention can be used to determine the specificity of the candidate modulator. Selectivity is evident, for example, in the field of chemotherapy, where the selectivity of a chemical to be toxic towards cancerous cells, but not towards non-cancerous cells, is obviously desirable. Selective modulators are preferable because they have fewer side effects in the clinical setting. The selectivity of a candidate modulator can be established *in vitro* by testing the toxicity and effect of a candidate modulator on a plurality of cell lines that exhibit a variety of cellular pathways and sensitivities. The data obtained from these *in vitro* toxicity studies can be extended animal model studies,

including human clinical trials, to determine toxicity, efficacy, and selectivity of the candidate modulator.

The identification of chemical, modulator, or therapeutic compositions.

5 The invention includes compositions, such as novel chemicals, and therapeutics identified by at least one method of the present invention as having activity as either a CYP450 substrate or inhibitor by the operation of methods, systems or components described herein. Novel chemicals, as used herein, do not include chemicals already publicly known in the art to be useful drugs or modulators as of the filing date of this application. Typically, a chemical would be identified as having CYP450 activity from
10 using the present invention and then its structure revealed from a proprietary database of chemical structures or determined using analytical techniques such as mass spectroscopy.

One embodiment of the invention is a chemical with useful activity, comprising a chemical identified by the method herein described. Such compositions include small organic molecules, nucleic acids, peptides and other molecules readily synthesized by
15 techniques available in the art and developed in the future. For example, the following combinatorial compounds are suitable for screening as candidate drugs: peptoids (PCT Publication No. WO 91/19735, 26 Dec. 1991), encoded peptides (PCT Publication No. WO 93/20242, 14 Oct. 1993), random bio-oligomers (PCT Publication WO 92/00091, 9 Jan. 1992), benzodiazepines (U.S. Pat. No. 5,288,514), diversomeres such as hydantoins.
20 benzodiazepines and dipeptides (Hobbs DeWitt. S. et al., Proc. Nat. Acad. Sci. USA 90: 6909-6913 (1993)), vinylogous polypeptides (Hagihara et al., J. Amer. Chem. Soc. 114: 6568 (1992)), nonpeptidal peptidomimetics with a Beta-D-Glucose scaffolding (Hirschmann, R. et al., J. Amer. Chem. Soc. 114: 9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen. C. et al., J. Amer. Chem. Soc. 116:2661
25 (1994)), oligocarbamates (Cho, C.Y. et al., Science 261: 1303 (1993)), and/or peptidyl phosphonates (Campbell, D.A. et al., J. Org. Chem. 59: 658 (1994)). See, generally, Gordon, E. M. et al., J. Med Chem. 37: 1385 (1994). The contents of all of the aforementioned publications are incorporated herein by reference.

The present invention also encompasses the compositions, identified by the methods
30 of the present invention, in a pharmaceutical compositions comprising a pharmaceutically acceptable carrier prepared for storage and subsequent administration, which have a pharmaceutically effective amount of the products disclosed above in a pharmaceutically

acceptable carrier or diluent. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A.R. Gennaro edit. 1985). Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. For example, sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid may be added as preservatives. In addition, antioxidants and suspending agents may be used.

These compositions may be formulated and used as tablets, capsules or elixirs for oral administration; suppositories for rectal administration; sterile solutions, suspensions for injectable administration; and the like. Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Suitable excipients are, for example, water, saline, dextrose, mannitol, lactose, lecithin, albumin, sodium glutamate, cysteine hydrochloride, and the like. In addition, if desired, the injectable pharmaceutical compositions may contain minor amounts of nontoxic auxiliary substances, such as wetting agents, pH buffering agents, and the like. If desired, absorption enhancing preparations (e.g., liposomes), may be utilized.

The pharmaceutically effective amount of the composition required as a dose will depend on the route of administration, the type of animal being treated, and the physical characteristics of the specific animal under consideration. The dose can be tailored to achieve a desired effect, but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

In practicing the methods of the invention, the products or compositions can be used alone or in combination with one another, or in combination with other therapeutic or diagnostic agents. These products can be utilized *in vivo*, ordinarily in a mammal, preferably in a human, or *in vitro*. In employing them *in vivo*, the products or compositions can be administered to the mammal in a variety of ways, including parenterally, intravenously, subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety of dosage forms. Such methods may also be applied to testing chemical activity *in vivo*.

As will be readily apparent to one skilled in the art, the useful *in vivo* dosage to be administered and the particular mode of administration will vary depending upon the age,

weight and mammalian species treated, the particular compounds employed, and the specific use for which these compounds are employed. The determination of effective dosage levels, that is the dosage levels necessary to achieve the desired result, can be accomplished by one skilled in the art using routine pharmacological methods. Typically, human clinical applications of products are commenced at lower dosage levels, with dosage level being increased until the desired effect is achieved. Alternatively, acceptable *in vitro* studies can be used to establish useful doses and routes of administration of the compositions identified by the present methods using established pharmacological methods.

In non-human animal studies, applications of potential products are commenced at higher dosage levels, with dosage being decreased until the desired effect is no longer achieved or adverse side effects disappear. The dosage for the products of the present invention can range broadly depending upon the desired affects and the therapeutic indication. Typically, dosages may be between about 10 microg/kg and 100 mg/kg body weight, preferably between about 100 microg/kg and 10 mg/kg body weight. Administration is preferably oral on a daily basis.

The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See e.g., Fingl *et al.*, in The Pharmacological Basis of Therapeutics, 1975. It should be noted that the attending physician would know how to and when to terminate, interrupt, or adjust administration due to toxicity, or to organ dysfunctions. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered dose in the management of the disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the dose and perhaps dose frequency, will also vary according to the age, body weight, and response of the individual patient. A program comparable to that discussed above may be used in veterinary medicine.

Depending on the specific conditions being treated, such agents may be formulated and administered systemically or locally. A variety of techniques for formulation and administration may be found in Remington's Pharmaceutical Sciences, 18th Ed., Mack Publishing Co., Easton, PA (1990). Suitable administration routes may include oral, rectal,

transdermal, vaginal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

5 For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For such transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art. Use of pharmaceutically acceptable carriers to formulate the compounds herein disclosed for the practice of the invention into dosages suitable for
10 systemic administration is within the scope of the invention. With proper choice of carrier and suitable manufacturing practice, the compositions of the present invention, in particular, those formulated as solutions, may be administered parenterally, such as by intravenous injection. The compounds can be formulated readily using pharmaceutically acceptable carriers well known in the art into dosages suitable for oral administration. Such
15 carriers enable the compounds of the invention to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

Agents intended to be administered intracellularly may be administered using techniques well known to those of ordinary skill in the art. For example, such agents may
20 be encapsulated into liposomes, then administered as described above. All molecules present in an aqueous solution at the time of liposome formation are incorporated into the aqueous interior. The liposomal contents are both protected from the external micro-environment and, because liposomes fuse with cell membranes, are efficiently delivered into the cell cytoplasm. Additionally, due to their hydrophobicity, small organic molecules
25 may be directly administered intracellularly.

Pharmaceutical compositions suitable for use as herein described include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. In
30 addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically.

The preparations formulated for oral administration may be in the form of tablets, dragees, capsules, or solutions. The pharmaceutical compositions of the present invention may be manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levitating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Pharmaceutical preparations for oral use can be obtained by combining the active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for

identification or to characterize different combinations of active compound doses. Such formulations can be made using methods known in the art (see, for example, U.S. Patent Nos. 5,733,888 (injectable compositions); 5,726,181 (poorly water soluble compounds); 5,707,641 (therapeutically active proteins or peptides); 5,667,809 (lipophilic agents); 5,576,012 (solubilizing polymeric agents); 5,707,615 (anti-viral formulations); 5,683,676 (particulate medicaments); 5,654,286 (topical formulations); 5,688,529 (oral suspensions); 5,445,829 (extended release formulations); 5,653,987 (liquid formulations); 5,641,515 (controlled release formulations) and 5,601,845 (spheroid formulations).

The following examples are meant to describe the inventors' preferred modes of carrying out the invention, *i.e.*, of preparing, characterizing, and using the preferred embodiments of the invention. Variations in the details of the particular methods employed and of the precise chemical compositions employed will undoubtedly be appreciated by those of skill in the art.

EXAMPLE 1

Synthesis of Fluorogenic Substrates for CYP450

Preparation of Benzyloxymethylresorufin (BOMR)

For all syntheses of the compounds of the invention, as herein described in this and the following EXAMPLES, the following protocols were (or are, with respect to EXAMPLE 7) followed unless so stated: Reaction conditions were (or are) carried out under atmospheric nitrogen. All solvents utilized were dried over 3Å molecular sieves. All chemicals and reagents were used as purchased without further purification unless stated. Benzychloromethylether was purchased from Fluka Chemie AG, Resorufin was purchased from Aldrich Chemical Co., 7-Hydroxy-3-trifluoromethycoumarin and 7-Hydroxy-3-cyanocoumarin were purchased from Molecular Probes and all were used as received. *See also* Wolfbeis, Otto, *Z. Naturforsch.* (1977) 32a, 1065-1067. Column chromatography was executed with J.T. Baker silica gel (particle size = 0.04-0.061 mm) using solvent combinations determined via initial TLC analysis with Merck Kieselgel 60 F₂₅₄, precoated silica gel plates. The ¹H NMR spectra, recorded at 500 MHz, were analyzed by NuMega Resonance Labs, Inc. Mass spectra were measured by ESI with a PE-SCIEX API 150EX.

Benzyloxymethylresorufin (BOMR) was prepared as follows: A suspension of resorufin, sodium salt, (235 mg, 1 mmol) and K₂CO₃ (248mg, 1.5mmol) in DMF (15 mL)

was vigorously stirred at 0-5 °C for 25 min. Benzylchloromethylether (2.32 mL, 10.0 mmol), was then added quickly to the reaction mixture. . The dark red mixture was stirred at 0 °C for 1.5 hrs. After which time the reaction turned to an orange solution. The reaction was monitored to completeness by TLC (R_f = 0.42, 1:1 EtOAc:Hex. and R_f = 0.05, CHCl_3).
5 The reaction was then brought up in Et_2O (35 mL), and extracted with saturated NaHCO_3 (30 mL). The aqueous layer was extracted two more times with Et_2O (30 mL). The etheral and the resorufin bilayer was then combined and filtered through celite. The filtrate was then dried with anhydrous NaSO_4 and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl_3) gave the pure
10 Benzyloxymethoxyresorufin as an orange solid (106 mg, 32%). ^1H NMR (500 MHz, CDCl_3): δ 4.74 (s, 2H), 5.39 (s, 2H), 6.32 (s, 1H), 6.85-6.83 (m, 1H), 7.06-7.09 (m, 2H), 7.30-7.37(m, 5H), 7.42 (d, 1H), 7.72(d, 1H).

EXAMPLE 2

15 Preparation of 7-Benzyloxymethoxy-3-cyanocoumarin (BOMCC)

7-Benzyloxymethoxy-3-cyanocoumarin (BOMCC) was prepared as follows:

A mixture of 7-Hydroxy-3-cyanocoumarin, (187 mg, 1 mmol) and K_2CO_3 (248 mg, 1.5 mmol), in DMF (15 mL) was vigorously stirred at 0°C for 25 min. Benzylchloromethylether (2.32 mL, 10.0 mmol), was then added quickly to the reaction.
20 The bright yellow mixture was stirred at 0 °C for 45 min. After which time the reaction turned to a colorless solution. The reaction was monitored to completeness by TLC (R_f = 0.5, 1:1 EtOAc:Hex. and R_f = 0.24, CHCl_3). The reaction was then brought up in Et_2O (35 mL), and extracted with saturated NaHCO_3 (30 mL). The aqueous layer was extracted two more times with Et_2O (30 mL). The etheral layer was then combined then dried with
25 anhydrous NaSO_4 and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl_3) gave the pure 7-Benzyloxymethoxy-3-cyanocoumarin as a white solid (9.21 mg, 3%). ^1H NMR (500 MHz, CDCl_3): δ 4.73 (s, 2H), 5.39 (s, 2H), 7.08 (m, 2H), 7.32 (m, 5H), 7.49 (d, 1H), 8.17 (s, 1H).

30

EXAMPLE 3

Preparation of 7-(*p*-methoxybenzyloxy-4-trifluorocoumarin (MOBFC)

7-(*p*-methoxybenzyloxy-4-trifluorocoumarin (MOBFC) was prepared as follows: A mixture of 7-Hydroxy-3-trifluoromethycoumarin, (230 mg, 1 mmol), K₂CO₃ (248 mg, 1.5 mmol), and KI (1.66 g, 10 mmol) in DMF (15 mL) was vigorously stirred at 25 °C for 25 min. Paramethoxybenzylchloride (1.35 mL, 10.0 mmol), was then added quickly to the reaction. The bright yellow mixture was stirred at 25 °C for 1 hr. After which time the reaction turned to a colorless solution. The reaction was monitored to completeness by TLC (R_f = 0.67, 1:1 EtOAc:Hex. and R_f = 0.3 CHCl₃). The reaction was then brought up in Et₂O (35 mL), and extracted with saturated NaHCO₃ (30 mL). The aqueous layer was extracted two more times with Et₂O (30 mL). The etheral layer were combined then dried with anhydrous NaSO₄ and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl₃) gave the pure 7-Paramethoxybenzyl-4-trifluorocoumarin as a white solid (280 mg, 80%). ¹H NMR (500 MHz, CDCl₃): δ 3.83 (s, 3H), 5.08 (s, 2H), 6.62 (s, 1H), 6.94 (m, 4H), 7.39 (m, 2H), 7.62 (m, 1H).

EXAMPLE 4

Preparation of Octyloxymethylresorufin (OOMR)

Octyloxymethylresorufin (OOMR) was prepared as follows: A suspension of resorufin, sodium salt, (235 mg, 1 mmol) and K₂CO₃ (248mg, 1.5 mmol) in DMF (15 mL) was vigorously stirred at 0-5 °C for 25 min. Bromomethyl octyl ether (2.20 mL, 10.0 mmol), was then added quickly to the reaction mixture. The reaction was stirred at 0-5 °C for 1.5 h during which time, the dark red reaction mixture turned to an orange solution. The reaction was allowed to continue to stir at 0-5 °C while monitoring by TLC (R_f = 0.44, 1:1 EtOAc:Hex. and R_f = 0.05, CHCl₃) and stopped at the time when product decomposition was detected. The reaction was then brought up in Et₂O (35 mL), extracted with 30mL of a saturated NaHCO₃ solution. The aqueous layer was extracted two more times with Et₂O (30 mL), the ether fractions were then combined and filtered through celite. The filtrate was then dried with anhydrous NaSO₄ and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl₃) yielded 62 mg of the purified octyloxy-methylresorufin (OOMR) as an orange solid. ¹H

NMR (500 MHz, CDCl_3): δ 8.3 (t, 3H), 1.21-1.31 (m, 12H), 3.68 (t, 2H), 5.31 (s, 1H), 6.32 (s, 1H), 6.83 (dd, 1H), 7.02-7.06 (m, 2H), 7.42 (d, 1H), 7.72 (d, 1H).

EXAMPLE 5

5 Preparation of 7-Methyloxymethyloxy-4-trifluorocoumarin (MOMFC)

7-Methyloxymethyloxy-4-trifluorocoumarin (MOMFC) was prepared as follows: A mixture of 7-hydroxy-4-trifluoromethylcoumarin (230 mg, 1 mmol) and K_2CO_3 (248 mg, 1.5 mmol), in DMF (15 mL) was vigorously stirred at 0-5 °C for 25 min. Bromomethyl methyl ether (0.97 mL, 10.0 mmol), was then added quickly to the reaction. The bright yellow mixture was stirred at 0-5 °C for 45 min during which time the reaction turned to a colorless solution. The reaction was allowed to continue to stir at 0-5 °C while monitoring by TLC (R_f = 0.54, 1:1 EtOAc:Hexane and R_f = 0.24, CHCl_3) and stopped at the time when product decomposition was detected. The reaction was then brought up in Et_2O (35 mL), extracted with 30 mL of a saturated NaHCO_3 solution. The aqueous layer was extracted two more times with Et_2O (30 mL), the ether fractions were then combined, dried with anhydrous NaSO_4 , filtered and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl_3) gave 11 mg of the purified 7-methyloxymethyloxy-4-trifluoromethylcoumarin (MOMFC) as a white solid. ^1H NMR (500 MHz, CDCl_3): δ 03.49 (s, 3H), 5.26 (s, 2H), 6.64 (s, 1H), 7.08-7.02 (m, 2H), 7.64 (d, 1H).

EXAMPLE 6

Preparation of Paramethoxybenzylresorufin (MOBR)





Paramethoxybenzylresorufin (MOBR) was prepared as follows: A mixture of resorufin, sodium salt, (235 mg, 1 mmol) and K_2CO_3 (248 mg, 1.5 mmol), in DMF (15 mL) was vigorously stirred at 0 °C for 25 min. Paramethoxybenzylchloride (1.35 mL, 10.0 mmol), was then added quickly to the reaction. The dark red mixture was stirred at 25 °C for 1.5 hrs. After which time the reaction turned to an orange solution. The reaction was monitored to completeness by TLC (R_f = 0.32, 1:1 EtOAc:Hex. and R_f = 0.05, CHCl_3). The reaction was then brought up in Et_2O (35 mL), and extracted with saturated NaHCO_3 (30 mL). The aqueous layer was extracted two more times with Et_2O (30 mL). The ethereal and the resorufin bilayer was then combined and filtered through celite. The filtrate was then

dried with anhydrous NaSO_4 and evaporated under reduced pressure. Chromatography of the crude product on silica gel (gradient 0-5% MeOH in CHCl_3) gave the pure Paramethoxybenzylresorufin as an orange solid (60 mg, 18%). ^1H NMR (500 MHz, CDCl_3): δ 3.83 (s, 3H), 5.10 (s, 2H), 6.32 (s, 1H), 6.82-6.88 (m, 2H), 6.94 (d, 2H), 6.99-7.01 (dd, 1H), 7.37 (d, 2H), 7.42 (d, 1H), 7.70(d, 1H).

EXAMPLE 7

Preparation of Other Optical CYP450 Sensors of the Invention


The following reaction schemes are used to synthesize other optical CYP450
 5 sensors of the present invention:



EXAMPLE 8

Kinetics of Benzyl-oxymethylresorufin (BOMR) Toward CYP 3A4

Applying the present invention to "modify" benzylresorufin (BR) leads, in one
 embodiment of the invention, to a compound of the invention, benzyl-oxymethylresorufin
 (BOMR), which has the following structure:



5

15

20

25

Detecting the Presence of CYP450 Inhibitors; Inhibition Assays

30

the plate, for final assay concentrations 1.3 mM NADP⁺, 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The drug inhibitors, miconazole, erythromycin, verapamil, diltiazem, ethinylestradiol, tamoxifen, and the substrates, digoxin, estrone, estradiol, warfarin, prednisone, and acetamidophen, were diluted from stock solutions of 10 mM in acetonitrile to 100 μ M in 100 mM K⁺ phosphate. 10 μ l of this dilution were added to appropriate wells on the plate for a final inhibitor concentration of 10 μ M. The CYP 3A4 was diluted to yield a solution containing 2 pmol/10ul in 100 mM K⁺ phosphate buffer, and 10 μ l was added to appropriate wells on plate. 20 μ l buffer was added to wells containing standards. The drug inhibitors were allowed to pre-incubated with the CYP 3A4 enzyme for 1 hr prior to the addition of the BOMR sensor. The BOMR sensor was diluted to 4 μ M (4X final assay concentration) in 100 mM K⁺ phosphate buffer, and 25 μ l was added to appropriate wells on the plate.

Data for a product fluorescence standard calibration curve was generated in the following manner. Resorufin was diluted to 40 μ M in K⁺ phosphate buffer, and seven consecutive 1:2 dilutions were made. 25 μ l of each dilution was added to the appropriate wells on the plate containing 75 μ l of 100 mM K⁺ phosphate, pH 8.0, and reading of the plates was begun immediately. For BOMR, the excitation filter was 530 nm and the emission filter was 580 nm.

The results of these experiments are shown in FIG. 5. Compounds known to be effective inhibitors (*e.g.*, miconazole and verapamil) inhibited the activity of CYP3A4 on BOMR by approximately 100%, effectively completely inhibiting the activity of CYP3A4 on BOMR.

Thus, as demonstrated with BOMR, and illustrated in FIG.s 4 and 5, introduction of the oxymethyl linker into fluorogenic substrates with long wavelength fluorophores, such as resorufin, yield the new CYP450 sensors of the invention with kinetic properties superior to those of the known, most closely-structurally related CYP450 substrates.

The utility in detecting drug-CYP450 interactions of selected oxymethyl-linker containing sensors of the present invention was further demonstrated for 7-benzyloxymethoxy-3-cyanocoumarin (BOMCC), and resorufin n-octyloxymethyl ether (OOMR), as illustrated in, respectively, FIG. 6, and FIG.s 6 and 7.

In FIG. 6, the results of a CYP 3A4 inhibition assay using BOMCC are illustrated. This FIG. illustrates that another compound of the invention, BOMCC, is useful as a means to detect the presence of inhibitors of the CYP450 enzyme CYP 3A4. This assay was performed in a 96-well plate at room temp and at a volume of 100 μ l/well. 1.82 X enzyme buffer was prepared and 55 μ l was added to each well on the plate, for final assay concentrations of 1.3 mM NADP⁺, 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The drug inhibitors, diphenylhydantoin, propranolol, imipramine, lansoprazole, pentamidine, and tranlycypromine, were diluted from stock solutions of 10 mM in acetonitrile to 100 μ M in 100 mM K⁺ phosphate. 10 μ l of this dilution was added to appropriate wells on the plate for a final inhibitor concentration of 10 μ M. The CYP3A4 was diluted to a yield a solution containing 2 pmol/10 μ l in 100 mM K⁺ phosphate buffer, and 10 μ l was added to appropriate wells on the plate. 20 μ l buffer was added to wells that contained standards. The drug inhibitors were allowed to pre-incubated with the 3A4 enzyme for 1 hour prior to the addition of the BOMCC substrate. The BOMCC sensor was diluted to 40 μ M (4X final assay concentration) in 100 mM K⁺ phosphate buffer, pH 8.0, and 25 μ l was added to appropriate wells.

Data for a product fluorescence standard calibration curve was generated in the following manner: 7-hydroxy-3-cyanocoumarin was diluted to 40 μ M in K⁺ phosphate buffer, and seven consecutive 1:2 dilutions were made. 25 μ l of each dilution was added to the appropriate wells on the plates containing 75 μ l of 100 mM K⁺ phosphate, pH 8.0, and immediately begin reading the plates. For BOMCC, the excitation filter was 395 nm and the emission filter was 460 nm.

In FIG. 7, the results of a CYP 2C9 inhibition assay using two compounds of the invention, BOMCC and OOMR, are illustrated. This FIG. illustrates that BOMCC and OOMR are useful as means to detect the presence of inhibitors of the CYP450 enzyme 2C9. This assay was performed in a 96-well plate at room temperature and at a volume of 100 μ l/well. 1.82 X enzyme buffer was prepared and 55 μ l was added to each well on the plate, for final assay concentrations 1.3 mM NADP⁺, 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The drug inhibitors, diclofenac, phenytoin, tenoxicam, tolbutamide, and sulfinpyrazone, were diluted from stock solutions of 10 mM in acetonitrile to 100 μ M in

100 mM K⁺ phosphate. 10 μ l of this dilution was added to appropriate wells on the plate for a final inhibitor concentration of 10 μ M. The CYP2C9 was diluted to a yield a solution containing 10 pmol/10 μ l in 100 mM K⁺ phosphate buffer, and 10 μ l was added to appropriate well on the plate. 20 μ l buffer was added to wells that contain standards. The
5 drug inhibitors were allowed to pre-incubated with the 2C9 enzyme for 1 hr prior to the addition of the BOMCC or OOMR sensor. The BOMCC sensor was diluted to 40 μ M (4X final assay concentration) in 100 mM K⁺ phosphate buffer, pH 8.0, and 25 μ l was added to the appropriate wells. The OOMR sensor was diluted to 8 μ M (4X final assay concentration) in 100 mM K⁺ phosphate buffer, and 25 μ l was added to appropriate wells
10 on the plate. The standards 7-hydroxy-3-cyanocoumarin and resorufin were diluted to 40 μ M in K⁺ phosphate buffer, to make seven 1:2 dilutions. 25 μ l of each dilution was added to the appropriate wells on the plate containing 75 μ l of 100 mM K⁺ phosphate, pH 8.0, and reading of the plates began immediately. For BOMCC, the excitation filter was 395 nm and the emission filter was 460 nm. For MOBR, the excitation filter was 530 nm and
15 the emission filter was 580 nm.

EXAMPLE 10









Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 3A4

A variety of the oxymethyl-containing sensors of the invention have been tested
20 against known human CYP450 isozymes predominantly involved with drug metabolism in humans. New substrates suitable for high throughput screening were found for CYP 3A4, CYP 2C19, CYP 2C9, CYP 1A2 and CYP 2B6. Tables 1 through 9 as described in detail in this and the following EXAMPLES, provide data regarding the kinetic properties of various fluorogenic CYP450 sensors of the present invention, as contrasted to the kinetic
25 properties of the most closely structurally-related, and currently-available, fluorogenic CYP450 substrates.

Table 1 has was prepared according to the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 1 correspond to specific fluorogenic substrates
30 tested against CYP 3A4; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. For further understanding these terms in the context of this invention, attention is

directed to FIG. 4 and the analysis of FIG. 4. As will be appreciated by those of skill in the art, the oxymethyl analogs of the present invention (BOMR, BOMFC, BOMCC, and EOMR) exhibited more efficient conversion to the same fluorescent product than each of the most closely structurally-related substrates (respectively, BR, BFC, BCC, and ER).
5 Indeed, in all cases presently studied, except for the case of one fluorogenic CYP450 substrate evaluated against one CYP450 enzyme (the effect of MOMFC as a substrate of CYP 2B6 as shown in Table 5), the oxymethyl derivatives of the present invention displayed improved kinetics over the most closely, structurally-related fluorogenic substrates.

Table 1.

3A4 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	type of kinetics
BR		0.08	0.02	0.1	5.3	377	Michaelis-Menten
BOMR		0.88	0.55	1.1	1.4	13095	Michaelis-Menten
BFC		0.81	0.04	n.a.	n.a.	n.a.	linear
BOMFC		1.41	0.11	4.0	22.0	3030	Michaelis-Menten
BCC		---	---	---	---	---	---
BOMCC		0.98	0.12	(10.0)	(89.0)	(1873)	sigmoidal
ER		---	---	---	---	---	---
EOMR		0.10	0.03	0.1	4.5	489	Michaelis-Menten







5 Kinetic properties of fluorogenic substrates with CYP 3A4. Benzylresorufin (BR) and 7-Benzyloxy-4-trifluoromethylcoumarin (BFC) are commercially available CYP450 substrates. Their oxymethyl analogs (BOMR, BOMFC) are more efficiently converted to the corresponding fluorescent product. The oxymethyl analogs of 7-Benzyloxy-3-cyanocoumarin and of ethylresorufin are better substrates than the parent substrates. (μ M = μ M; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)

EXAMPLE 11

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 2C19

Table 2 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 2 correspond to specific fluorogenic substrates tested against CYP 2C19; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. As will be appreciated by those of skill in the art, the oxymethyl analogs of the present invention (EOMCC, BOMCC, and MOMCC) exhibited more efficient conversion to the same fluorescent product than each of the most closely, structurally-related substrates (respectively, 3CEC, BCC, and 3CMC).

Table. 2.

2C19 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
3CEC		0.34	0.05	1.7	41.0	671	Michaelis-Menten
EOMCC		2.22	0.34	5.7	16.6	5723	Michaelis-Menten
BCC		0.06	0.02	0.1	6.4	234	Michaelis-Menten
BOMCC		0.68	0.16	1.9	14.0	2262	Michaelis-Menten
3CMC		0.07	0.01	n.a.	n.a.	n.a.	linear
MOMCC		0.38	0.14	0.5	3.2	2604	Michaelis-Menten

Kinetic properties of fluorogenic substrates with CYP2 C19. 3-Cyano-7-ethoxycoumarin (3CEC) is a commercially available CYP450 substrate. Its oxymethyl analog (EOMCC) is more efficiently converted to the corresponding fluorescent product. The oxymethyl analogs of 7-benzyloxy-3-cyanocoumarin and of 3-cyano-7-methoxycoumarin are better substrates than the parent substrates. (uM = μ M; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)







EXAMPLE 12

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 2C9

Table 3 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 3 correspond to specific fluorogenic substrates tested against CYP 2C19; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. As will be appreciated by those of skill in the art, the oxymethyl analogs of the present invention (MOMFC, BOMCC, and MOMCC) exhibited more efficient conversion to the same fluorescent product than each of the most closely, structurally-related substrates (respectively, MFC, BCC, and 3CMC).

15

Table 3.

2C9 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MFC		0.03	0.01	0.4	103.0	70	Michaelis-Menten
MOMFC		0.07	0.00	0.1	14.9	157	Michaelis-Menten
BCC		0.00	0.00	---	---	---	---
BOMCC		0.42	0.04	2.1	43.0	814	Michaelis-Menten
3CMC		0.04	0.00	0.4	70.9	85	linear
MOMCC		0.07	0.01	0.2	20	167	Michaelis-Menten



Kinetic properties of fluorogenic substrates with CYP 2C9. 7-Methoxy-4-trifluoromethylcoumarin (MFC) is a commercially available CYP450 substrate. Its oxymethyl analog (MOMFC) is more efficiently converted to the corresponding fluorescent product. The oxymethyl analogs of 7-benzyloxy-3-cyanocoumarin and of 3-cyano-7-methoxycoumarin are better substrates than the parent substrates. (uM = μ M; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)

EXAMPLE 13

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 1A2

Table 4 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 3 correspond to specific fluorogenic substrates tested against CYP 1A2; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. As will be appreciated by those of skill in the art, the oxymethyl analog of the present invention (EOMCC) exhibited more efficient conversion to the same fluorescent product than each of the most closely, structurally-related substrates (3CEC).

Table 4.

1A2 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	type of kinetics
3CEC		12.55	1.73	44.0	26.0	28205	Michaelis-Menten
EOMCC		9.95	1.77	21.0	11.0	31818	Michaelis-Menten

Kinetic properties of fluorogenic substrates with CYP 1A2. 3-Cyano-7-ethoxycoumarin (3CEC) is a commercially available CYP450 substrate. Its oxymethyl analog (EOMCC) is converted to the corresponding fluorescent product a little bit more efficiently (greater k_{cat}/K_m).







EXAMPLE 14

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 2B6

Table 5 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 5 correspond to specific fluorogenic substrates tested against CYP 2C19; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. As will be appreciated by those of skill in the art, two of the oxymethyl analogs of the present invention test in this EXAMPLE (BOMCC and BOMR) exhibited more efficient conversion to the same fluorescent product than each of the most closely, structurally-related substrates (respectively, BCC, and BR).

As noted above, the case of MOMFC as a substrate of CYP 2B6 is the sole case in which the fluorogenic CYP450 substrate of the present invention did not exhibit improved kinetics, *i.e.*, more efficient conversion to the same fluorescent product, than the most closely, structurally-related substrate. in that case MFC. By the method used to identify this sole case, or comparable methods for selecting fluorogenic CYP450 substrate and CYP450 enzyme pairs, those of skill in the art may distinguish the most desirable fluorogenic CYP450 substrates of the present invention for their particular use.

Table 5.

2B6 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	type of kinetics
MFC		2.53	1.63	2.9	1.4	34524	Michaelis-Menten
MOMFC		0.71	0.54	n.a.	n.a.	n.a.	not Michaelis-Menten
BCC		0.07	0.04	0.1	1.3	1064	Michaelis-Menten
BOMCC		3.05	0.38	66.0	52.0	21154	Michaelis-Menten
BR		0.42	0.42	n.a.	n.a.	n.a.	not Michaelis-Menten
BOMR		0.71	0.28	0.8	0.73	18265	Michaelis-Menten

5 Kinetic properties of fluorogenic substrates with CYP 2B6. 7-Methoxy-4-trifluoromethylcoumarin (MFC) is a commercially available CYP450 substrate. This is the occasion on which the oxymethyl analog (MOMFC) is less efficiently converted to the corresponding fluorescent product found to date (11/15/98). The oxymethyl analogs of 7-benzyloxy-3-cyanocoumarin and of benzylresorufin are better substrates than the parent

10 substrates. (μ M = μ M; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)










EXAMPLE 15

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 3A4 and CYP 2D6

Table 6 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 6 correspond to specific fluorogenic substrates tested against CYP 3A4 and CYP 2D6, as indicated; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected.

10

Table 6.

3A4 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M
OOMR		0.601	0.323	0.66	1.4	7857
BOM-DDAO		4.2041	0.4402	6	7	14286
OOMCC		0.961	0.227444	1.19	4.4	4508
MOMR		0.0338	0.0083	0.06	6	167
BRCBE		0.6241	0.1992	0.83	2.7	5123
MOBFC		0.3453	0.0214	0.7	13	897
2D6 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M
MOMR		0.21457	0.03396	0.34	8	708
<i>MOBR</i>		0.0409	0.0185	0.049	3.1	263
<i>IPCC</i>		0.0378	0.0076	0.1	18	93

Other substrates first synthesized and tested on CYP 3A4 and CYP 2D6. Oxymethyl ether derivatives of the invention (OOMR, BOM-DDAO, OOMCC, MOMR;) are listed in bold; other ethers are listed in italics.









EXAMPLE 16

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 2C9 and CYP 2C19

Table 7 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 7 correspond to specific fluorogenic substrates tested against CYP 2C9 and CYP 2C19, as indicated; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected.

10

Table 7.

2C9 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M
OOMR		0.31	0.137	0.4	2.6	2564
<i>MOBFC</i>		<i>0.188</i>	<i>0.02</i>	<i>0.29</i>	<i>7.7</i>	<i>628</i>
2C19 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M
OOMCC		1.043	0.179	1.7	6.9	4106
OOMR		0.13	0.1	0.17	0.7	4048
MOMR		0.062	0.06	<i>n.d.</i>	<i>n.d.</i>	<i>n.d.</i>
<i>MOBR</i>		<i>0.69</i>	<i>1.26</i>	<i>n.d.</i>	<i>n.d.</i>	<i>n.d.</i>
<i>DMMC</i>		<i>0.23</i>	<i>0.03</i>	<i>0.4</i>	<i>8.5</i>	<i>784</i>
IPCC		0.232	0.027	1.95	57.2	568







5 Other substrates first synthesized and tested on CYP 2C9 and CYP 2C19. Oxymethyl ether derivatives of the invention (OOMR, OOMCC, MOMR) are listed in bold; other ethers are listed italics.

EXAMPLE 17

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 3A4 and CYP 2D6

Table 8 was prepared according to the same general methodology of Table 1, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 8 correspond to specific fluorogenic substrates tested against CYP 3A4 and CYP 2D6, as indicated; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics detected. As will be appreciated by those of skill in the art, the oxyphenylmethyl analogs of the present invention (MOBFC, MOBR) exhibited more efficient conversion to the same fluorescent product than each of the most closely, structurally-related substrates (respectively, MFC and MR). Indeed, in all cases presently studied, the oxyphenylmethyl derivatives of the present invention displayed improved kinetics over the most closely, structurally-related fluorogenic substrates.

Table 8.

3A4 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MFC		---	---	---	---	---	---
MOBFC		0.35	0.02	0.7	13.0	897	Michaelis-Menten
2D6 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MFC		0.10	0.03	0.3	10.0	417	Michaelis-Menten
MOBFC		0.49	0.09	0.6	5.0	2067	Michaelis-Menten
MR		---	---	---	---	---	---
MOBR		0.04	0.02	0.0	3.1	263	Michaelis-Menten

Kinetic properties of fluorogenic substrates with CYP 3A4 and CYP 2D6. 7-Methoxy-4-trifluoromethylcoumarin (MFC) and Methylresorufin (MR) are commercially available CYP450 substrates. Their oxyphenylmethyl analogs (MOBFC and MOBR) are more efficiently converted to the corresponding fluorescent products. (uM = μ M; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)

EXAMPLE 18

Analysis of the Relative Kinetics of Fluorogenic Substrates of CYP 2C9, CYP 2C19 And
2B6

5 Table 9 was prepared according to the same general methodology of Table 1, the
general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*,
"Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 9
correspond to specific fluorogenic substrates tested against CYP 2C9 and CYP 2C19 and
CYP 2B6, as indicated; the columns correspond to, respectively, the abbreviations of the
10 fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at
1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics
detected. As will be appreciated by those of skill in the art, the oxyphenylmethyl analogs
of the present invention (MOBFC, MOBR) exhibited more efficient conversion to the same
fluorescent product than each of the most closely, structurally-related substrates
(respectively, MFC and MR). Indeed, in all cases presently studied, the oxyphenylmethyl
15 derivatives of the present invention displayed improved kinetics over the most closely,
structurally-related fluorogenic substrates.

Table 9.

2C9 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MFC		0.03	not done	0.4	103.0	70	Michaelis- Menten
MOBFC		0.19	0.02	0.3	7.7	628	Michaelis- Menten
MR		---	---	---	---	---	---
MOBR		0.02	0.02	0.0	0.4	952	Michaelis- Menten
2C19 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MR		---	---	---	---	---	---
MOBR		0.69	1.26	n.a.	n.a.	n.a.	not MMK
2B6 Abbr.	Substrate Structure	v (10uM) (min-1)	v (1.25uM) (min-1)	kcat (min-1)	Km (microM)	kcat/Km s-1/M	type of kinetics
MR		---	---	---	---	---	---
MOBR		0.10	0.08	0.4	0.1	58333	Michaelis- Menten

Kinetic properties of fluorogenic substrates with CYP 2C9, CYP 2C19 and CYP 2B6. 7-Methoxy-4-trifluoromethylcoumarin (MFC) and Methylresorufin (MR) are commercially available CYP450 substrates. Their oxyphenylmethyl analogs (MOBFC and MOBR) are more efficiently converted to the corresponding fluorescent products. (uM = μM ; n.a. = not applicable; --- too low to quantify, () from Michaelis-Menten fit)

EXAMPLE 19

Determining the Apparent Inhibition Constants (k_i) of Drugs that Interact
with CYP450 2D6

To demonstrate the effectiveness of MOBFC as a sensor for a specific subfamily of human CYP450, CYP 2D6 and its use in determining apparent inhibition constants following experiment was conducted. CYP 2D6 was incubated with 10 μ M concentrations of various known inhibitors and drug substrates and residual CYP 2D6 activity assayed with the fluorogenic substrate MOBFC. The assay was performed in a 96-well plate at room temp and at a volume of 100 μ l/well. 4X enzyme buffer was prepared and 25 μ l was added to each well on the plate. for final assay concentrations of 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The drug inhibitors, quinidine, chlorpheniramine, yohimbine, imipramine, amjaline, propranolol, doxorubicin, haloperidol and corynanthine were diluted from stock solutions of 10 mM in acetonitrile to 120 μ M in 100 mM K⁺ phosphate. Six consecutive 1:3 dilutions were made and 50 μ l of each was added to appropriate wells on the assay plate. The CYP2D6 was diluted to a solution of 2 pmol/10 μ l in 100 mM K⁺ phosphate and 10 μ l was added to each well. 20 μ l of buffer was added to each well containing standard. The drug inhibitors were allowed to pre-incubated with the CYP 2D6 enzyme for 1 hr prior to the addition of the MOBFC substrate. The MOBFC substrate was diluted to 26.6 μ M (6.7X final assay concentration) in 100 mM K⁺ phosphate buffer, and 15 μ l was added to appropriate wells on the plate. Data for a product fluorescence standard calibration curve was generated in the following manner. Hydroxy-trifluoromethylcoumarin was diluted to 100 μ M in K⁺ phosphate buffer, and seven consecutive 1:2 dilutions were made. 10 μ l of each dilution was added to the appropriate wells on the plate containing 90 μ l of 100 mM K⁺ phosphate, pH 8.0. After addition of 10 μ L of 13 mM NADP⁺ solutions to all wells the assay plate was placed into the fluorescence microtiter plate reader and fluorescence measured at 3 minute intervals for 60 minutes. For MOBFC, the excitation filter was 395/25 nm and the emission filter was 530/25 nm. IC₅₀ values (value for 50% inhibition of fluorogenic substrate turnover) were determined and converted to apparent k_i values according to the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993).

Table 10.

DRUG	Apparent K_i values [μ M]
Quinidine	0.15
Chlorpheniramine	2.5
Yohimbine	10
Imipramine	0.1
Amjaline	> 30
Propanolol	> 30
Doxorubicin	8
Haloperidol	3
Corynanthine	> 30

Apparent k_i values for inhibition of CYP 2D6 by drugs known to interact with the enzyme determined from IC_{50} values of inhibition of MOBFC metabolism by the enzyme.

EXAMPLE 20

Preparation of 7-Benzylloxymethyloxycoumarin-3-carboxylic acid succinimidyl ester

7-Benzylloxymethyloxycoumarin-3-carboxylic acid succinimidyl ester was prepared by following procedure: A mixture of 7-Hydroxycoumarin-3-carboxylic acid succinimidyl ester, (303 mg, 1 mmol) and dry potassium carbonate (248 mg, 1.5 mmol), in dry dimethylformamide (15 mL) was vigorously stirred at 0 °C for 25 min. Benzylchloromethylether (2.32 mL, 10.0 mmol), was then added quickly to the reaction. The bright yellow mixture was stirred at 0 °C for 45 min. and for 2 hrs. at 25 °C. After which time the reaction turned to a colorless solution. The reaction was monitored by TLC (R_f = 0.5, 1:1 EtOAc:Hex. and R_f = 0.24, $CHCl_3$). After of the coumarin starting material the reaction medium was diluted with diethylether (100 mL) and extracted with 50 mL of 5% aqueous acetic acid. The ether layer was separated and dried over anhydrous sodium sulfate, filtered and the solvents evaporated under reduced pressure. The solid was recrystallized from methanol and washed with hexanes (20 mL, 0 °C). The product, -


Benzyloxymethoxycoumarin-3-carboxylic acid succinimidyl ester, was dried under reduced pressure yielding a white solid (85 mg, 20%). ¹H NMR (500 MHz, CDCl₃): 2.91 (s, 4H), 4.74 (s, 2H), 5.40 (s, 2H), 7.07 (m, 2H), 7.32 (m, 5H), 7.53 (d, 1H), 8.75 (s, 1H).

5

EXAMPLE 21

Coupling of 7-benzyloxymethoxycoumarin-3-carboxylic acid succinimidyl ester with racemic 1,2-diaminocyclohexanes) to give the product designated BOM-09B

Five (5) μmoles benzyloxymethoxycoumarin-3-carboxylic acid succinimidyl ester in dry dimethylformamide (50 μL) were mixed with a 1 M solution a racemic mixture of 1,2-diaminocyclohexanes (50 μmoles) in a plastic centrifuge tube. The reaction was allowed to proceed with sonication at room temperature for 2 hrs, after which time 500 μL of deionized water was added to the tube and a white precipitate formed. The reaction was then spun down in a centrifuge and the solvent decanted. Results from UV-Vis spectra (absorbance maximum at 340nm) and electrospray MS (M+H = 423, M+Na = 445) performed on a sample of the solid were consistent with the following product structure (mixture of stereo isomers):



BOM-09B

EXAMPLE 22

Determination of the rate of hydroxylation of BOM-09B by CYP 3A4

The racemic compound mixture (designated BOM-09B) was tested for activity with cytochrome P450 isozymes. BOM-09B showed particularly high activity with the CYP 3A4 isozyme. The CYP 3A4 assay was performed in a 96-well plate at 37 °C in a volume of 100 µL/well. BOM-09B was diluted from a stock solution of 1 mM in acetonitrile to a 4X concentration of 80 µM in 100 mM K⁺ phosphate buffer of which 25 µl was added to the appropriate wells. Enzyme buffer was prepared and 65µl was added to each well on the plate, for final assay concentrations of 1.3 mM NADP⁺, 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The cytochrome P450 isozyme CYP3A4 was diluted to give a 2 pmol enzyme per well. Enzymic conversion of the substrate to products was allowed to proceed for 1 hour with fluorescence reads taken every 4 minutes on a fluorescence microtiter plate reader. The solution was illuminated with an 395/25 nm excitation filter and fluorescence emission was detected through a 460/40 nm the emission filter. The rate of conversion of this substrate (BOM-09B) was compared with the substrate BOMCC under identical conditions and found to be half that of BOMCC (conversion rate _(BOM-09B) = 0.75pmol substrate/ pmol enzyme · min).


20

EXAMPLE 23

Synthesis of Fluorogenic Substrate Libraries from Highly Fluorescent Phenolic Dyes

Libraries of ethers of 7-hydroxycoumarins and resorufin derivatives are synthesized as outlined below, *i.e.*, the reaction paths leading to libraries of fluorogenic CYP450 candidate substrates are shown below. 7-Hydroxycoumarin-3-carboxylic acid succinimidyl ester is commercially available from Molecular Probes. The resorufin starting materials are readily prepared by following the procedures of U.S. Patent Nos. 4,954,630 and 5,304,645, which describe the preparation of the acids and their conversion to the active esters using TSTU. The active esters of the dyes are stable to alkylation conditions needed to prepare ethers of the dye phenols. After alkylation the resulting fluorogenic dye ethers are modified at the active ester moiety by reaction with a library of primary and secondary aliphatic amines. The aliphatic sidechains are chosen to include diverse aromatic and heterocyclic moieties. 20 diamines are also included in the amine library. When reacted

with the active dye esters in large molar excess, the reactions with diamines result in positively charged candidate substrates, which are screened for activity against the CYP 2D6 isozyme, which are known to prefer positively charged substrates.



5

Compounds are purified by column chromatography or recrystallization after the alkylation of the dye phenol. Coupling with the amines is performed on a 10 μ mol scale in dimethylformamide and usually proceeds in high yield. The reactions are followed in parallel by thin layer chromatography to ensure completion of the reactions and to detect and remove poorly reacting library members. The solvent is removed in high vacuum and excess amine removed by suspending the residue in 10% aqueous acetic acid, followed by recovery of the product by centrifugation. This procedure leads to compounds with sufficient purity (tested by TLC) for initial testing for metabolism by CYP450 enzymes. Promising substrates are resynthesized on a larger scale (100 μ mol), purified by chromatography including separation of regio-isomers (resorufin-based substrates) and analyzed by electro-spray MS and analyzed by 1 H-NMR.

15

EXAMPLE 24

Testing Libraries of Fluorogenic Substrates Against a Set of Human CYP450 Isozymes

The libraries of newly synthesized putative substrates are dissolved at 2 mM concentration in appropriate water miscible organic solvents, with acetonitrile generally preferred, because up to 2% is generally tolerated by CYP450 enzymes. The different ether derivatives of any phenolic dye have similar extinction coefficients, allowing for calibration of the substrate concentration by absorbance. The solutions are transferred to 96 well storage plates to allow multiple automated parallel dispensation into 96 well microtitre assay plates. For rapid testing, all assays are performed in microtitre plates, using a Fluorstar or Cytofluor fluorescence plate reader to obtain the enzyme rates.

Initial experiments determine the conditions for each CYP450 isozyme (using commercially available human CYP450 isozymes expressed in insect microsomes from GENTEST) giving linear rates of product formation; the rate is proportional to the concentration of enzyme. The CYP450 isozymes tested to find more active substrates are: CYP 3A4, CYP 2D6, CYP 2C9, CYP 2C19. Testing also includes the CYP450 isozymes CYP 1A2, CYP 2E1, CYP 2B6, and CYP 2A6. This is to assess whether any substrate that is active with one of the isozymes 3A4, 2D6, or 2C9 and 2C19 is selective for that isozyme. Each isozyme requires slightly different conditions, and optimized variables include pH, NADPH concentration, concentration of CYP450, whether it is necessary to add cytochrome b as a cofactor, time of incubation, and effect of temperature, and other variables that will be apparent to those of skill in the art. In the initial screen for substrates, coumarin-based candidates are tested at 5 and 20 μ M concentrations, resorufin-based candidates at 2 and 10 μ M. The choice of different concentrations for ethers of resorufins versus ethers of coumarins takes into account the lower aqueous solubility of resorufin derivatives compared to coumarin derivatives and our finding in preliminary experiments that resorufin derivatives, being more hydrophobic, tend to bind more avidly to CYP450 enzymes (lower K_m). One pmol insect-expressed enzyme coexpressed with NADPH-cytochrome P450 reductase per well are used. The assay buffer will contain 10mM Mg^{2+} and the appropriate ionic strength of the assay solution is adjusted with from concentrated buffer stock solutions. NADPH needed for NADPH-cytochrome P450 reductase is supplied in the form of 1.3 mM $NADP^+$, which is converted to steady levels of NADPH by added glucose-6-phosphate dehydrogenase and 3mM glucose-6-phosphate in the assay

buffer. Apparent k_{cat} and K_m values for all active candidates are determined from eight-point dilutions of each substrate in duplicate, using the results from the preliminary tests to determine the actual concentration range for the accurate kinetic evaluation.

Optimization of assay conditions is accomplished through statistically validated Design of Experiments methodologies, using the commercial software package "Design-Expert[®]" produced by Stat-Ease[®] Inc. The data shown in FIG.s 4 and 5 were obtained with initial optimization.

EXAMPLE 25

Directed Synthesis of Fluorogenic Substrate Sets

Substrates found in the initial round of synthesis and testing are resynthesized on a larger scale (100 μ mol) and purified by chromatography and/or recrystallization. Resorufin regio-isomeric ethers obtained in the synthesis are separated and kinetic properties determined for each separate isomer, as described above. Kinetic data obtained for these substrates will be used to direct the synthesis of a few small focused libraries. Additional alkyl halides and amines, closely related to the ones that result in activity with the isozyme, are purchased or synthesized with the goal of obtaining substrates with even higher activity and substrates that may be isozyme specific. The same synthetic routes as discussed in EXAMPLE 23, are followed, except that all compounds are purified and analyzed by NMR and MS before performing enzyme kinetics. These substrate candidates are tested in duplicate in eight-point dilutions using the results from the structurally-related substrates to determine the actual concentration range for the accurate kinetic evaluation.

EXAMPLE 26

Validation of Isozyme-Specific Substrates Using Human Liver Microsomes

Because human liver microsomes contain a range of CYP450 isozymes, only substrates that are specific for one of the insect-expressed human CYP450 enzymes are tested on commercially available human liver microsomal preparations. This verifies that, as generally expected, the specificity seen with the insect microsomal CYP450s is maintained in human liver microsomes. Initially, conditions for the assays are those specified by the suppliers of the microsomes. However, because the new substrates may have different kinetics to those for which the published conditions were designed, some

optimization as described in EXAMPLE 24 is performed. All assays are carried out in 96 or 384-well microtitre plates as in EXAMPLE 24.

Specificity for one isozyme in the human liver microsomal preparations is confirmed by a lack of metabolism of the substrate in the presence of a selective CYP450 isozyme inhibitor for the isozyme being investigated. For example, inhibitors selective for CYP 3A4, CYP 2D6, and CYP 2C9 are troleandomycin, quinidine, and sulfaphenazole respectively. In addition, the fluorogenic substrates/sensors are used to determine IC_{50} values for a panel of known CYP450 isozyme inhibitors, and the data compared to published values. For this step, an 8-point concentration curve of the inhibitors is performed in duplicate. Some difficulty may be encountered in that the published literature contains a large range of IC_{50} values for any given inhibitor, often because of different experimental conditions between studies. Our results will be compared with the more relevant published studies, which have the most similar assay conditions.

EXAMPLE 27

Validate Screens Against Known CYP450 Inhibitors and Substrates

The most relevant CYP450 isozymes (CYP3A4, CYP2D6, CYP2C9, CYP2C19) are screened with their most appropriate novel fluorogenic substrates/sensors against a library of compounds containing known CYP 450 inhibitors. For this EXAMPLE, GENTEST recombinant human CYP450 isozymes expressed in insect microsomes are used. The library to be tested is the generic pharmacophore library from Microsource®, which has 480 biologically active molecules, including known CYP450 inhibitors and substrates.

Since the ultimate commercial value of the new CYP450 substrates/sensors is realized if the assays are adapted to high throughput, automated screening protocols, it is necessary to verify that the assay conditions developed in EXAMPLE 24 are suitable for automated use, and if not, to modify them appropriately. This is common practice with any assay which is to be run on a robotic system, and involves checking such parameters as: enzyme and substrate stability to allow a large number of assays to be run without constant manual intervention; reproducibility of the assay with the automated liquid handling systems; specific incubation times and temperatures suitable for the robotic scheduler; appropriate data capture and reduction routines, and other like parameters. It is

also determined whether a "pre-read" of the plates prior to adding the NADPH to initiate the reaction will be necessary, since this can sometimes eliminate false positives caused by fluorescent compounds in the library.

Initially all compounds are tested at 10 μ M concentration using the optimized robotic conditions, to determine in a simple hit/miss mode which compounds are interacting with which CYP450 isozyme. Although this involves testing four enzymes against 480 compounds (approx. 2,000 assays with controls), this only requires 25 96-well or 6 384-well microtitre plates, which can be run in a single day using currently available automated formats.

All hits are retested at 1 and 10 μ M, and tested at 10 μ M using a redox-sensitive red fluorescent dye identified to be suitable for checking that a compound is not interfering with the cytochrome P450 reductase step. IC₅₀ values are determined (using an eight-point curve in duplicate) for the known inhibitors or substrates and compare the data to published values. For the new substrates/sensors to be deemed suitable for routine high throughput screening of compounds as part of the drug development process, the assays must detect 100% of compounds with affinities for the relevant CYP450 isozyme of <1 μ M, and >90% of compounds with affinities between 1 and 10 μ M.

EXAMPLE 28

Determination of whether a test compound is a substrate for a CYP450 isozyme

To determine whether a test compound is a substrate for a CYP450 isozyme the following experiment is conducted. A preparation of human CYP450 isozyme is treated with test compound for an incubation period of several hours under conditions suitable for metabolism of the test compound by the CYP450 isozyme. The residual CYP450 isozyme activity is assayed with a fluorogenic substrate for that CYP450 isozyme. The CYP450 isozyme is also treated with the same test compound for the same period of time but in the absence of NADP⁺, a condition that does not allow test compound metabolism. Following the incubation period, NADP⁺ is added, and the residual CYP450 isozyme activity is assayed with a fluorogenic substrate for that CYP450 isozyme. A CYP450 isozyme activity assayed under conditions suitable for metabolism of the test compound that is higher than the activity of the enzyme under conditions that do not allow test compound

metabolism during the incubation period indicates that the test compound is a substrate of the CYP450 isozyme.

The assay is performed in a 96-well plate at room temperature and at a volume of 100 μ l/well. 4X enzyme buffer is prepared and 25 μ l is added to each well on the plate, for
5 final assay concentrations of 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM $MgCl_2$ in a K^+ phosphate buffer of suitable concentration and at pH 8.0. The test compound is dissolved to 20 μ M concentration in water and 50 μ L of the solution is added to two wells each, followed by addition of 10 μ L of buffer containing 10pmol of the CYP450 isozyme. One of the two wells now receives
10 10 μ L of 13 mM NADP $^+$ and the test compound in both wells is incubated with the CYP450 isozyme for 2 hrs. Following incubation, the other well receives 10 μ L of 13 mM NADP $^+$ and both receive fluorogenic substrate, suitable for detection of activity of the CYP450 isozyme, in a 5 μ L volume of buffer. The microtiter assay plate is transferred into the fluorescence microtiter plate reader and well fluorescence is measured at 3-minute
15 intervals for 60 minutes. The rate in increase of well fluorescence is used to assess residual CYP450 isozyme activity in the wells. A result in which the residual CYP450 activity in the well that receives NADP $^+$ prior to the incubation with the test compound is higher than in the duplicate well to which NADP $^+$ is added after the incubation period indicates that the test compound is a substrate for the CYP450 isozyme.

20

EXAMPLE 29

Analysis of fluorescence signal from enzymatic metabolism of substrate over background signal from reagent additions

To demonstrate the superior optical qualities of the molecular sensors of the
25 invention, one of which is improved signal over background, side by side comparisons with currently-available fluorogenic substrates were performed. The assays were performed in a 96-well plate at room temperature and at a volume of 100 μ l/well. 2X NADPH/Recycling buffer was added to the plate at a volume of 50 μ l for final assay concentrations of 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM $MgCl_2$
30 in 100 mM K^+ phosphate buffer, pH 8.0 (with exception of the CYP2C9 and CYP2C19 assays for which 50 mM K^+ phosphate, pH 8.0 were used). (See Table 11.) The enzyme was made up in the appropriate K^+ phosphate buffer at 10X and 10 μ l/well was added to the

appropriate wells. (See Table 11.) The plate was read at 3 minute intervals for 12 minutes to obtain a background fluorescence levels. Readings were briefly interrupted to allow for the addition of 10 μ l of 10X NADP+ in K+ phosphate buffer to each well. (See Table 11.) The readings were resumed at 3 minute intervals for another 12 minutes. Again the

5 readings were interrupted to allow for the addition of 3.3X substrate at a volume of 30 μ l/well (See substrate concentrations in table 11.). Enzymatic conversion of the substrate to products was allowed to proceed for 1 hour with fluorescence reads taken at 3 or 4 minute intervals on a fluorescence microtiter plate reader at the appropriate excitation and emission wavelength (listed for each substrate in Table 11). FIGS. 8, 9, 10, 11 and 12

10 illustrate the fluorescence intensity changes, or lack thereof, for addition of reagents and substrate to the wells and the signal from enzymic metabolism of the substrate. To permit quantitative comparison between sensors that were or would be metabolized to yield fluorescent products of differing optical properties, the signal intensity corresponding to each well was normalized by division with the initial signal intensity (wells containing

15 enzyme and buffer). Figures 8, 9, 10, 11 and 12 illustrate the improved signal over background of oxymethyl and oxyphenylmethyl linker containing substrates of this invention (solid traces) over prior commercially available substrates (broken lines). The cumulative background signal resulting from the addition of NADP+ and from addition of the substrate manifest itself by a fluorescence signal greater than one, 3 minutes following

20 the substrate addition (second arrow indicates time of substrate addition). The later increase in signal, approximately four minutes after the second addition and at later time points, is believed to be due to the enzymatic conversion of the substrate to the product. In Figures 8, 9, 10, 11 and 12, enzymatic signal over reagent addition background, as a measure of the performance of the assay, is greater for the oxymethyl-linker containing

25 sensors of this invention (solid traces) than for prior commercially available substrates (broken lines). Figure 8 illustrates the superior signal to background of BOMR (solid trace) versus benzylresorufin (broken trace) with the CYP3A4 isozyme. Figure 9 illustrates the superior signal to background of BOMCC (solid trace) versus 7-Benzyloxy-4-trifluoromethylcoumarin (BFC, broken trace) with the CYP3A4 isozyme. Figure 10

30 illustrates the superior signal to background of MOBFC (solid trace) versus AMMC (Gentest) (broken trace) with the CYP2D6 isozyme. Figure 11 illustrates the superior signal to background of both OOMR (solid trace) and BOMCC (solid trace) versus 7-

Methoxy-4-trifluoromethylcoumarin (MFC, broken trace) with the CYP2C9 isozyme. Figure 12 illustrates the superior signal to background of EOMCC (solid trace) versus 3-cyano-7-ethoxycoumarin (CEC, broken trace) with the CYP2C19 isozyme.

Table 11.

Isozyme	CYP3A4	CYP3A4	CYP2D6	CYP2C9	CYP2C19
Substrates in separate wells each	1 μ M BOMR 5 μ M Benzyl-resorufin	20 μ M BOMCC 20 μ M <i>BFC</i>	5 μ M MOBFC 2.5 μ M <i>AMMC</i>	2.5 μ M OOMR 10 μ M BOMCC 40 μ M <i>MFC</i>	20 μ M EOMCC 20 μ M CEC
Glucose-6-phosphate (mM)	33	33	33	33	33
Glucose-6-phosphate dehydrogenase (units/ml)	0.4	0.4	0.4	0.4	0.4
MgCl ₂ (mM)	10	10	10	10	10
K ⁺ phosphate buffer, pH 8.0 (mM)	100	100	100	50	50
Enzyme concentration (pmol/100 μ l)	0.5	0.5	2	1	0.5
NADP ⁺ (μ M)	100	100	30	100	100
Excitation/Emission (nm)	Ex. 530(25)/ Em. 605(50)	Ex. 405(20)/ Em. 460(40) <i>Ex. 405(40)/ Em. 490(40)</i>	Ex. 405(40)/ Em. 490(40) <i>Ex. 360(40)/ Em. 460(40)</i>	Ex. 530(25)/ Em. 605(50) Ex. 405(20)/ Em. 460(40) <i>Ex. 405(40)/ Em. 490(40)</i>	Ex. 405(20)/ Em. 460(40)
Center wavelength (bandwidth)					
Corresponding Figures	FIG. 8	FIG. 9.	FIG. 10.	FIG. 11	FIG. 12

5 CYP450 assay condition for each CYP450 isozyme. Molarities indicate concentrations after addition of all reagents to each well. Substrates were compared in side-by-side wells at the indicated concentrations (second row). Different font types for substrates indicate the matching filters in second to last row (e.g., filter set in italics was used for substrate in italics).

EXAMPLE 30

Preparation of Dibenzylloxymethylfluorescein (DBOMF)

Dibenzylloxymethylfluorescein was prepared as follows:

5 A mixture of fluorescein (332 mg, 1 mmol) and diisopropylethylamine (870 μ L, 5 mmol), in chloroform (15 mL) was vigorously stirred at 0 °C for 25 minutes. Benzylchloromethylether (2.32 mL, 10.0 mmol) was then added quickly to the reaction. The reaction gave off gas upon the addition, and was allowed to stir for 2 hours at 25 °C. The reaction was monitored by TLC (R_f = 0.36, 5%MeOH : 95% CHCl_3). The reaction solution was then purified by chromatography on silica gel (gradient 0-5% MeOH in CHCl_3 which gave
10 the pure DBOMF as an orange solid (154 mg, 27%). ^1H NMR (500 MHz, CDCl_3): 4.37 (q, 2H), 4.70 (s, 2H), 5.29 (q, 2H), 5.33 (s, 2H), 6.47 (d, 1H), 6.57 (m, 1H), 6.86-6.93 (m, 3H), 7.14-7.24 (m, 2H), 7.25-7.36 (m, 10H), 7.67-7.77 (m, 2H), 8.23 (d, 1H).

15

EXAMPLE 31

Preparation of Benzyloxymethylfluorescein (BOMF)

Benzyloxymethylfluorescein (BOMF) was prepared as follows:




20 A mixture of fluorescein (332mg, 1 mmol) and diisopropylethylamine (870 μ L, 5 mmol), in chloroform (15 mL) was vigorously stirred at 0 °C for 25 minutes. Benzylchloromethylether (2.32 mL, 10.0 mmol), was then added quickly to the reaction. The reaction gave off gas upon the addition, and was allowed to stir for 2 hours at 25 °C. The reaction was monitored by TLC (R_f = 0.56, 5%MeOH : 95% CHCl_3). The reaction solution was then purified by chromatography on silica gel (gradient 0-5% MeOH in CHCl_3 which gave
25 the pure BOMF as an orange solid (36 mg, 8%). ^1H NMR (500 MHz, CDCl_3): 4.72 (s, 2H), 5.31 (s, 2H), 6.52 (q, 1H), 6.59 (d, 1H), 6.68-6.76 (m, 3H), 7.00 (d, 1H), 7.17 (d, 1H), 7.29-7.37 (m, 5H), 7.60-7.69 (m, 2H), 8.02 (d, 1H).

EXAMPLE 32

Analysis of the Relative Kinetics of Fluorescein-based Fluorogenic Substrates with CYP3A4

Table 12 was prepared according to the same general methodology of Table 1, that is, the general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*, "Enzyme Assays," Oxford University Press, 277-313 (1993). The rows of Table 12 correspond to specific fluorescein-based fluorogenic substrates tested against CYP 3A4; the columns correspond to, respectively, the abbreviations of the fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at 1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, v_{max} (for sigmoidal kinetics), $K_{1/2max}$ (concentration at 1/2 maximal velocity for sigmoidal kinetics) and the types of kinetics detected. Dibenzylfluorescein (DBF) was purchased from Gentest, Woburn, Mass. The CYP3A4 assay was performed in a 96-well plate at room temperature in a volume of 100 μ L/well. DBF, BOMF and DBOMF were diluted from a stock solution of 2 mM in acetonitrile to a 4X concentration of 80 μ M in 100 mM K⁺ phosphate buffer. Seven 1:2 dilutions were made of each substrate of which 25 μ l was added to the appropriate wells. 1.54X Enzyme buffer was prepared and 65 μ l was added to each well on the plate. for final assay concentrations of 100 μ M NADP⁺, 3.3 mM glucose-6-phosphate, 0.4 units/ml glucose-6-phosphate dehydrogenase and 10 mM MgCl₂ in 100 mM K⁺ phosphate, pH 8.0. The cytochrome P450 isozyme CYP3A4 was diluted to give a 0.5 pmol enzyme per well and added in a volume of 10 μ l/well in K⁺ phosphate, pH 8.0. Enzymatic conversion of the substrate to products was allowed to proceed for 1 hour with fluorescence reads taken every four minutes on a fluorescence microtiter plate reader. The solution was illuminated using an 485/20 nm excitation filter and fluorescence emission was detected through a 530/25 nm the emission filter. As will be appreciated by those of skill in the art, the oxymethyl analog sensors of the present invention (DBOMF and BOMF) exhibited more efficient conversion to the same fluorescent product than the structurally related sensor (DBF).

Table 12.

3A4 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	v_{max} (min ⁻¹)	$K_{1/2max}$ (microM)	type of kinetics
DBF		4.40	3.27	n.a.	n.a.	n.a.	4.30	0.80	sigmoidal
DBOMF		15.50	6.00	n.a.	n.a.	n.a.	16.00	1.50	sigmoidal
BOMF		3.40	0.53	9.0	15.0	10000	n.a.	n.a.	Michaelis Menten

Kinetic properties of fluorogenic fluorescein-based substrates with CYP 3A4.

Dibenzylfluorescein (DBF) was purchased from Gentest. Its oxyphenylmethyl analogue

- 5 Dibenzylloxymethylfluorescein (DBOMF) is more efficiently converted to the corresponding fluorescent product. Benzyloxymethylfluorescein (BOMF) is another structurally related oxyphenylmethyl ether derivative of fluorescein which is also efficiently converted to fluorescein and displays Michaelis Menten type kinetics.

EXAMPLE 33

Analysis of the Relative Kinetics of Fluorescein-based Fluorogenic Substrates with
CYP 2C9

5 Table 13 was prepared according to the same general methodology of Table 12 in
Example 32, that is the general method described in Henderson, P.J.F., *Statistical Analysis*
of Enzyme Kinetic Data, "Enzyme Assays," Oxford University Press, 277-313 (1993). In
deviation from the protocol described in Example 32, the phosphate-based buffer (100mM
K⁺ phosphate, pH 8.0) was replaced throughout the procedure by 200mM Tris-HCl buffer,
10 pH 7.5. The rows of Table 13 correspond to specific fluorescein-based fluorogenic sensors
tested against CYP 2C9; the columns correspond to, respectively, the abbreviations of the
fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at
1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics
detected. As will be appreciated by those of skill in the art, the mono-oxymethyl analog of
15 fluorescein of the present invention, BOMF, exhibited very efficient conversion to the
fluorescent product compared to substrates listed in Table 3.


2C9 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	type of kinetics
BOMF		2.84	1.28	4.5	4.5	16666	Michaelis Menten

Table 13.

Kinetic properties of substrate BOMF with CYP 2C9.

EXAMPLE 34

Analysis of the Relative Kinetics of Fluorescein-based Fluorogenic Substrates with
CYP450 isozyme 2C8

Table 14 was prepared according to the same general methodology of Table 12, the
5 general method described in Henderson, P.J.F., *Statistical Analysis of Enzyme Kinetic Data*,
"Enzyme Assays," Oxford University Press, 277-313 (1993). In deviation from the
protocol described in Example 32, the phosphate-based buffer (100mM K⁺ phosphate, pH
8.0) was replaced throughout the procedure by 100mM K⁺ phosphate buffer, pH 7.5. The
rows of Table 13 correspond to specific fluorescein-based fluorogenic sensors tested
10 against CYP 2C8; the columns correspond to, respectively, the abbreviations of the
fluorogenic substrate, the chemical structure, the turnover rate at 10 μ M, the turnover rate at
1.25 μ M, k_{cat} , and K_m values, the ratio of k_{cat} and K_m values, and the types of kinetics
detected. Specifically, the substrate dibenzylfluorescein (DBF) (purchased from Gentest,
Woburn, Mass.) and the sensor of the present invention DBOMF were tested for activity
15 with CYP 2C8. As will be appreciated by those of skill in the art, the oxymethyl analog of
the sensor of the present invention (DBOMF) exhibited more efficient conversion to the
same fluorescent product than the structurally-related substrate (DBF).



2C8 Abbr.	Substrate Structure	v (10 μ M) (min ⁻¹)	v (1.25 μ M) (min ⁻¹)	k_{cat} (min ⁻¹)	K_m (microM)	k_{cat}/K_m s ⁻¹ /M	type of kinetics
DBF		0.37	0.16	0.4	2.0	3333	Michaelis Menten
DBOMF		1.54	0.55	2.1	3.6	9722	Michaelis Menten

Table 14.

Kinetic properties of fluorescein-based fluorogenic substrates with CYP 2C8.

Dibenzylfluorescein (DBF) was purchased from Gentest. Its oxyphenylmethyl analogue

- 5 Dibenzylloxymethylfluorescein (DBOMF) is more efficiently converted to the corresponding fluorescent product.

EXAMPLE 35

Procedure for Screening of Compounds for Interactions with CYP 3A4

A general assay procedure for CYP 3A4 isozyme inhibition assays using sensors of the present invention were performed and is described. Tables 15 and 16 describe the specific buffer conditions used in assaying inhibition of this enzyme. Assays were run in 96-well black-walled clear-bottom plates, 100 μ l/well at room temperature. Fluorescence is measured on a fluorescence microtiter plate reader.

BOMR was dissolved to 2 mM concentration in acetonitrile via the addition of 500 μ l acetonitrile to a vial containing 1 μ mol substrate. BOMCC was dissolved in acetonitrile to make 10 mM stock solutions by addition of 100 μ l acetonitrile to a vial containing 1 μ mol substrate. Sodium resorufin dye standard was dissolved to 1 mM in distilled water; 3-cyano-7-hydroxycoumarin dye standard was prepared at 1mM in DMSO. All aqueous solution were prepared at the beginning of the experiment and kept on ice until use.

1. One nanomole test compound or inhibitor was dispensed into the plate well in distilled water (40 μ l of 25 μ M compound in distilled water), for a final assay concentration of compound of 10 μ M.

Alternatively, compounds are dispensed from an organic solution and dried on the plate, in which case 40 μ l/well distilled water is added and the plate incubated at 37°C for 1 hr to aid re-dissolution of compounds. Control wells receive the appropriate inhibition controls for each isozyme. Also, inhibitors are added to wells that contain the dye standards to permit addition of substrate and enzyme to all wells of the plate.

2. 20 μ l/well of 300 mM K⁺ Phosphate buffer, pH 8.0, was added.
3. 30 μ l/well Enzyme/Recycling System buffer (Table 15) was added. The dye standards were added to the appropriate wells on the plate. 10 μ l/well of dye standards were added for final assay concentrations of 5, 2.5 and 1.25 μ M.

Table 15.

	BOMR	BOMCC
K ⁺ Phosphate, pH 8.0 (mM)	100	100
Glucose-6-phosphate (mM)	11	11
Glucose-6-phosphate dehydrogenase (units/ml)	1.33	1.33
MgCl ₂ (mM)	33.33	33.33
Enzyme (nanoM)	16.5	16.5
Equivalent to pmol/well	0.5	0.5

Concentration of ingredients in stock of Enzyme/Recycling System Buffer (3.3X)

- 5 4. The plate was incubated for 20 minutes at room temperature to aid interaction of compounds with CYP3A4 (in the absence of enzyme turnover). A pre-reading of the plate was taken (See Table 16 below for excitation and emission filters for each substrate).
- 10 5. Ten μ l/well of Substrate/NADP⁺ buffer in 100 mM K⁺ phosphate, pH 8.0 (Table 16) was added. Immediately thereafter, the plate was transferred into the fluorescence microtiter plate reader for kinetic analyses, or alternatively, the plate is allowed to sit in the dark for the appropriate reaction time and an endpoint reading is taken at the appropriate time.

15 Table 16.

Substrate Name	BOMR	BOMCC
Substrate (μ M)	50	200
NADP ⁺ . (mM)	1	1
Assay time (minutes)	45	45
Excitation, center (width)/	530(25)/ 605(55)	405(20)/ 460(40)
Emission, center (width)		

Substrate/NADP⁺ Buffer (10X)

EXAMPLE 36

Screening of 160 Compounds for Interactions with CYP 3A4

Following the general procedure outlined in Example 35, 160 compounds purchased from Chembridge, San Diego, Calif., were screened for interactions with CYP 3A4. One nanomole of compound was dispensed per well, eighty per 96 well plate. Into the remaining 16 wells was dispensed the following: Three nmol miconazole as control for 100% inhibition, a moderate inhibitor verapamil at four concentrations to indicate sensitivity of the assay and a fluorescence product standard (resorufin sodium salt). The results of applying the procedure outlined in Example 35 with 5 μ M BOMR as substrate are illustrated in FIG. 13. The potency of the compounds in the sample (expressed in % inhibition) compared to the miconazole control was graphed for all compounds. Several compounds with intermediate potency (30-60% inhibition relative to control) were detected. Eight compounds in the sampling displayed inhibitory activity greater 60%. As will be appreciated by those skilled in the art, these results demonstrate that the oxymethyl linker-containing sensors of the present invention are useful in, for example, detecting CYP 3A4 inhibitory activity in a variety and/or array of compounds.

EXAMPLE 37

Detection of Compound Substructures that Correlate with CYP 3A4 Inhibitory Activity

Following the general procedure outlined in Example 35, 160 compounds purchased from Chembridge were screened for interactions with CYP 3A4. Eight compounds in the sample displayed inhibitory activity greater 60%. The structures of six of these, and their percentage inhibitory activity compared to control, are illustrated in Fig. 14. As will be appreciated by those skilled in the art, the oxymethyl linker containing substrates are useful to detect compounds with related substructures as well as structurally unrelated compounds that display CYP 3A4 inhibitory activity. For example, a sensor of the present invention revealed inhibitory activity by two compounds containing planar aromatic and thiourea substructures (FIG. 14, left panel), and identified potent inhibitors having iodonium substructures (FIG. 14, center panel). Also, two other, less closely structurally-related compounds with aniline and/or heterocyclic motifs were identified as having inhibitory activity.

EXAMPLE 38

Process for Screening of Compounds for Interaction with CYP 2D6

5 A general assay procedure for CYP 2D6 isozyme inhibition assay using oxyphenylmethyl linker containing sensors of this invention was performed and is described. Tables 17 and 18 list the specific buffer conditions used with this enzyme. Assays were run in 96-well black-walled clear-bottom plates, 100 μ l/well at room temp. Fluorescence was measured on a fluorescence microtiter plate reader.

10 MOBFC substrate was dissolved in acetonitrile to make 10 mM stock solutions by addition of 100 μ l acetonitrile to a vial containing 1 μ mol substrate. Substrate stock solutions were stable when stored at 4°C in the dark. 7-Hydroxy-4trifluoromethylcoumarin dye standard was prepared at 1mM in DMSO. All aqueous solution were prepared at the beginning of the experiment and kept on ice until use.

- 15 1. One nanomole of test compound or inhibitor was dispensed into the plate well in distilled water (40 μ l of 25 μ M compound in distilled water), for a final assay concentration of compound of 10 μ M.

Alternatively, compounds are dispensed from an organic solution and dried on the plate, in which case 40 μ l/well distilled water is added and the plate is incubated at 37°C for 1 hour to aid re-dissolution of compounds. Control wells receive the appropriate inhibition controls for each isozyme. Also, inhibitors are added to wells that contain the dye standards to permit addition of substrate and enzyme to all wells of the plate.

- 20 2. 20 μ l/well of 300 mM K⁺ Phosphate buffer, pH 8.0 were added.
3. 30 μ l/well Enzyme/Recycling System buffer was added. The dye standards were added to the appropriate wells on the plate. 10 μ l/well of dye standards are added for final assay concentrations of 5, 2.5 and 1.25 μ M.

Table 17

	2D6
K ⁺ Phosphate, pH 8.0 (mM)	100
Glucose-6-phosphate (mM)	11
Glucose-6-phosphate dehydrogenase (units/ml)	1.33
MgCl ₂ (mM)	33.33
Enzyme (nM)	66
Equivalent to pmol/well	2

Concentration of ingredients in stock of Enzyme/Recycling System Buffer (3.3X)

- 5 4. Plates were incubated for 20 minutes at room temperature to aid the interaction of compounds with isozymes (in the absence of enzyme turnover). A pre-reading of the plate was taken.
5. 10 μ l/well of Substrate/NADP⁺ buffer in 100 mM K⁺ phosphate, pH 8.0 was added. The plate was immediately transfer into the fluorescence microtiter plate reader for kinetic analyses, or alternatively, the plate is allowed to sit in the dark for the appropriate reaction time (Table 18) and an endpoint reading is taken at the appropriate time.

Table 18.

15

	2D6
Substrate Name	MOBFC
Substrate (μ M)	50
NADP ⁺ , (mM)	0.3
Assay time (minutes)	120
Excitation, center (width)/	405(40)/
Emission, center (width)	490(40)

Substrate/NADP⁺ Buffer (10X)

EXAMPLE 39

Screening of 240 Compounds for Interactions with CYP 2D6

Following the general procedure outlined in Example 38, 240 compounds purchased from Chembridge (including the compounds screened in Example 36) were screened for interactions with CYP 2D6. One nanomole compound was dispensed per well, eighty per 96 well plate. Into the remaining 16 wells was dispensed: 1 nmol quinidine as control for 100% inhibition and a fluorescence product standard (7-hydroxy-4-trifluoromethylcoumarin). The results of applying the procedure outlined in Example 38 with 5 μ M MOBFC as substrate are illustrated in FIG. 15. The potency of the compounds in the sampling expressed in % inhibition compared to quinidine control was graphed for all compounds. Several compounds with intermediate potency (30-60% inhibition relative to control) were detected. About 10% of compounds in the sample displayed inhibitory activity greater 60%. As will be appreciated by those skilled in the art, these results demonstrates that the oxyphenylmethyl linker containing sensors of the present invention are useful in, for example, detecting CYP 2D6 inhibitory activity in a variety and/or array of compounds.

EXAMPLE 40

Detection of Compound Substructures that Correlate with CYP 2D6 Inhibitory Activity

Following the general procedure outlined in Example 38, 240 compounds purchased from Chembridge were screened for interactions with CYP 2D6. Approximately ten percent of the compounds in the sampling array displayed inhibitory activity greater 60%. The structures of seven of these, and their percentage inhibitory activity compared to control, are illustrated in Fig. 16. As will be appreciated by those skilled in the art, the oxyphenylmethyl linker-containing sensors of the present invention are useful in, for example, detecting compounds with related substructures as well as in detecting structurally-unrelated compounds that display CYP 2D6 inhibitory activity. For example, a sensor of the present invention revealed the inhibitory activity of three structurally-related compounds that contain orthophenylenediamine substructures (FIG. 16, left panel), and revealed potent inhibitors having iodonium substructures (FIG. 16, center panel). Also, three other less closely structurally-related compounds, each containing residues that are positively charged at neutral pH, were determined to have inhibitory activity.

EXAMPLE 41

Method of Detection Whether The Inhibitory Activity Of A Compound Against A CYP4505 Isozyme Is Dependent On Its Metabolism or On Activation By The Enzyme

CYP450 enzymes are inhibited by compounds that interfere with substrate binding, the binding of molecular oxygen, and/or with the catalytic step in which the substrate is oxidized. See Ortiz de Montellano, P.R., Correia, M.A., Inhibition of cytochrome P450 enzymes. In: Cytochrome P450: Structure, mechanism and biochemistry, 2nd edition, 10 Plenum Press, New York (1995), 305-364. Compounds that can bind to the heme moiety in the active site in its ferric or ferrous state inhibit the enzyme; compounds that have affinity to additional structural motifs in the enzyme's active site being particularly potent. For example, compounds that contain imidazole and pyridine functionalities may bind the heme iron. Additionally, a variety of sulfur compounds, and olefin-containing and 15 acetylene-containing derivatives, inhibit CYP450 enzymes by being activated to species that may covalently bind to the enzymes active site. Also, certain amine-containing compounds are metabolized to intermediates that strongly bind the ferrous heme.

The CYP450 sensors of this invention have been used to assess whether the potency of CYP450 inhibition by a compound or drug candidate is dependent on NADPH- 20 dependent turnover (metabolism) of the compound. The procedure employed makes use of the observation that an inhibitor that is activated by the enzyme to an intermediate that irreversibly binds to the active site of the enzyme, such as a so-called suicide inhibitor, progressively inhibits the enzyme in a time dependent fashion. This observation is in contrast to a true competitive inhibitor whose potency varies little (or may possibly drop if it is metabolized) after the compound is pre-incubated with the CYP450 enzyme under 25 conditions that allow for turnover of the enzyme. A compound that is metabolized to a potent inhibitor or whose turnover "knocks out" the enzyme (non-competitive, suicide inhibitor), will appear more potent if pre-incubated with the enzyme under conditions that permit metabolism of such compound, as inhibition is metabolism-dependent and therefore 30 progressive with time. As will be appreciated by persons of skill in the art, there may exist compounds that may act both as suicide inhibitors that cause progressive inhibition and as competitive inhibitors of the enzyme. Compounds of this invention can be used to assess

whether a compound inhibits a CYP450 isozyme in a metabolism-dependent and time-dependent fashion.

Described below are the experimental assay procedures employed for the CYP 3A4 isozyme. These procedures permit assessment of progressive metabolism-dependent inhibition of the enzyme using sensors of this invention. Tables 19-22 list the specific buffer conditions used with this enzyme. Assays were run in 96-well black-walled clear-bottom plates, 100 μ l/well at room temperature. Fluorescence was measured on a fluorescence microtiter plate reader.

BOMR substrate was dissolved to 2 mM concentration in acetonitrile (add 500 μ l acetonitrile to vial containing 1 μ mol substrate). BOMCC substrate was dissolved in acetonitrile to make 10 mM stock solutions by addition of 100 μ l acetonitrile to vial containing 1 μ mol substrate. DBOMF substrate was dissolved in acetonitrile to make 2mM stock solution by addition of 500 μ l acetonitrile to vial containing 1 μ mol substrate. Sodium resorufin dye standard was dissolved to 1mM in distilled water, 3-Cyano-7-hydroxycoumarin dye standard was made up at 1mM in DMSO. Fluorescein dye standard was made up at 1mM in DMSO. All aqueous solution were made up fresh at the beginning of the experiment and kept on ice until use.

The assays were performed under two conditions, Conditions A (steps A1-A5) and Conditions B (Steps B1-B5). The results are summarized in Table 23.

Experimental conditions, Conditions A:

One nanomole of test compound or inhibitor was dispensed into the plate well in distilled water (40 μ l of 25 μ M compound in distilled water), for a final assay concentration of compound of 10 μ M.

6. 20 μ l/well of 300 mM K⁺ Phosphate buffer, pH 8.0 was added.

7. 30 μ l/well Enzyme/Recycling System buffer (See Table 19) was added. The dye standards were added to the appropriate wells on the plate. 10 μ l/well of dye standards were added for final assay concentrations of 5, 2.5 and 1.25 μ M.

Table 19.

	BOMR	BOMCC	DBOMF
K ⁺ Phosphate, pH 8.0 (mM)	100	100	100
Glucose-6-phosphate (mM)	11	11	11
Glucose-6-phosphate dehydrogenase (units/ml)	1.33	1.33	1.33
MgCl ₂ (mM)	33.33	33.33	33.33
Enzyme (nanomM)	16.5	16.5	16.5
Equivalent to pmol/well	0.5	0.5	0.5

Concentration of ingredients in stock of Enzyme/Recycling System Buffer (3.3X)

8. Plates were incubated for 20 minutes at room temperature to aid interaction of compounds with CYP3A4 (in the absence of enzyme turnover). A pre-reading of the plate was taken. (See Table 20 for excitation and emission filters used for each substrate).
9. 10 μ l/well of Substrate/NADP⁺ buffer in 100 mM K⁺ phosphate, pH 8.0 (Table 20) was added. Plates were transferred into a fluorescence microtiter plate reader for kinetic analyses.

Table 20.

Substrate Name	BOMR	BOMCC	DBOMF
Substrate (μ M)	50	200	20
NADP ⁺ , (mM)	1	1	1
Assay time (minutes)	45	45	45
Excitation, center (width)/ Emission, center (width)	530(25)/ 605(55)	405(20)/ 460(40)	485(20)/ 530(25)

Substrate/NADP⁺ Buffer (10X)

Experimental conditions, Conditions B:

- One nanomole of test compound or inhibitor was dispensed into the plate well in distilled water (40 μ l of 25 μ M compound in distilled water), for a final assay concentration of compound of 10 μ M.

B1. 20 μ l/well of 300 mM K⁺ Phosphate buffer, pH 8.0 was added.

- B2. 30 μ l/well Enzyme/Recycling System buffer (See Table 21) was added. The dye standards were added to the appropriate wells on the plate. 10 μ l/well of dye standards were added for final assay concentrations of 5, 2.5 and 1.25 μ M.

Table 21.

	BOMR	BOMCC	DBOMF
K ⁺ Phosphate, pH 8.0 (mM)	100	100	100
Glucose-6-phosphate (mM)	11	11	11
Glucose-6-phosphate dehydrogenase (units/ml)	1.33	1.33	1.33
NADP ⁺ (mM)	0.33	0.33	0.33
MgCl ₂ (mM)	33.33	33.33	33.33
Enzyme (nanoM)	16.5	16.5	16.5
Equivalent to pmol/well	0.5	0.5	0.5

Concentration of ingredients in stock of Enzyme/Recycling System Buffer (3.3X)

B3. Plates were incubated for 40 minutes at room temperature in the presence of NADP⁺ to permit interaction of compounds with CYP3A4 under conditions allowing for enzyme turnover. Pre-readings of the plates were taken. (See Table 22 for excitation and emission filters used for each substrate.)

B4. 10 μ l/well of Substrate buffer in 100 mM K⁺ phosphate, pH 8.0 (Table 22) was added. Plate were transferred into fluorescence microtiter plate reader for kinetic analyses.

Table 22.

Substrate Name	BOMR	BOMCC	DBOMF
Substrate (μ M)	50	200	20
Assay time (minutes)	20	20	20
Excitation, center (width)/ Emission, center (width)	530(25)/ 605(55)	405(20)/ 460(40)	485(20)/ 530(25)

Substrate/NADP⁺ Buffer (10X)

Data were collected for fluorescence intensities in each well containing 1 nanomole of drugs as listed in Table 23. Data were converted to reflect percent inhibition relative to the control. Controls were 10 μ M clotrimazole for Condition A, and 10 μ M ellipticine for Condition B. The results are presented in Table 23, in which percent inhibition of BOMR, BOMCC and DBOMF metabolism under both Conditions A and Conditions B, are listed for 15 drugs. Several candidate drugs (the nine top listed drugs) were more potent inhibitors of the metabolism of sensors of this invention under conditions permitting metabolism of the compound (condition B), than under condition A. Some large differences in percent inhibition values for drugs tested under the separate conditions are indicated in bold type (Table 23). DBOMF was particularly insensitive to inhibition by drugs under Condition A, but reliably indicated metabolism-dependent inhibition of the

same (Condition B). Data for DBOMF indicated that the top nine candidate drugs (Dicyclomine, Verapamil, Ellipticine, Erythromycin, Clemastine, Amiodarone, Mifepristone, Doxorubicin, Papaverine) possesses at least partial metabolism-dependent inhibitory activity against the CYP3A4 isozyme. As will be appreciated by persons skilled in the art, sensors of this invention are useful to assess whether a chemical, test compound, drug candidate or drug acts as a turnover-dependent inhibitor of a CYP450 isozyme. Moreover, compounds characterized as suicide inhibitors or suicide substrates, chemical CYP450 knock-out agents, or non-competitive CYP450 inhibitors may display turnover-dependent inhibition of a CYP450 isozyme and may be detected using sensors of this invention. Test compounds that are converted to metabolites with CYP450 inhibitory activity may also be detected (Product inhibitors).

Table 23.

SUBSTRATE	BOMR		BOMCC		DBOMF	
	A	B	A	B	A	B
Dicyclomine	77	97	76	98	30	86
Verapamil	83	95	88	101	36	89
Ellipticine	97	100	92	100	40	100
Erythromycin	90	99	93	98	30	85
Clemastine	62	85	82	95	32	80
Amiodarone	35	71	73	98	1	77
Mifepristone	35	81	93	98	22	98
Doxorubicin	29	72	81	79	3	33
Papaverine	69	92	34	99	48	90
Ticlopidine	77	84	13	55	-9	3
Isoniazid	-8	30	59	48	-9	3
Nitrofurantoin	-19	27	44	33	-9	-3
Diphenylhydantoin	-27	24	-11	21	21	5
Oxprenolol	-5	13	7	21	1	3
Ketoconazole	89	97	91	97	98	99

CYP3A4 inhibitory activity of 15 drugs under conditions A and B as assessed by sensors of this invention (BOMR, BOMCC, DBOMF). Numbers are given as % inhibition as compared to 100% inhibition control.

The various articles of the scientific and/or medical literature, and the U.S. and foreign patents and patent applications cited herein are hereby incorporated by reference; each constitutes a part of the disclosure of this specification. Furthermore, while specific embodiments, working examples, and prophetic examples of the invention have been described in detail to illustrate the broad applicability and principles underlying the

invention, it will be understood by those of skill in the art that the invention may be embodied otherwise without departing from such broad applicability and principles.

WE CLAIM:


1. A compound useful as a sensor for cytochrome P450 activity having the structure:

Y-L-Q, wherein:

5 Y is selected from the group consisting of Q as herein defined, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl; wherein if Y is selected from
 10 Q as herein defined, L is L', wherein L' is selected from the group of -(CR⁴H)(-OCR²H)_p, -(CR⁴H)(-O(substituted ortho-phenyl)CR²H)_p-, -(CR⁴H)(-O(substituted meta-phenyl)CR²H)_p-, and -(CR⁴H)(-O(substituted para-phenyl)CR²H)_p, wherein each R² and each R⁴ is separately selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl and p is a positive integer no
 15 greater than twelve;

20 L has the chemical structure L', as herein defined, or (-OCR²H)_p-, wherein each R² is separately selected from the group consisting of a hydrogen atom, saturated C₁-C₂₀ alkyl, unsaturated C₁-C₂₀ alkenyl, unsaturated C₁-C₂₀ alkynyl, substituted saturated C₁-C₂₀ alkyl, substituted unsaturated C₁-C₂₀ alkenyl, substituted unsaturated C₁-C₂₀ alkynyl, C₁-C₂₀ cycloalkyl, C₁-C₂₀ cycloalkenyl, substituted saturated C₁-C₂₀ cycloalkyl, substituted unsaturated C₁-C₂₀ cycloalkenyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, and p is a positive integer no
 25 greater than twelve; and


30 Q has a structure selected from the group consisting the following structures:




I




II




III




IV




V




VI




VII




VIII




IX




X




XI



XII




XIII




XIV


5




XV




XVI




XVII



XVIII




XIX




XX


5




XXI




XXII




XXIII



XXIV




XXV




XXVI

10



XXVII



XXVIII

wherein:

m is a positive integer no greater than five;

$R_a, R_b, R_c, R_d, R_e, R_f, R_g, R_h, R_i, R_j, R_k,$ and R_l are each separately selected from the group consisting of a hydrogen atom, a halogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano, nitro, azido, - SR_s , - OR_o , - $NR_{n1}R_{n2}$, - $N^+R_{n1}R_{n2}R_{n3}$, - $P^+R_{n1}R_{n2}R_{n3}$, - COR_c , - $C(=NOR_o)R_c$, - CSR_c , - $OCOR_c$, - $OCONR_{n1}R_{n2}$, - OCO_2R_c , - $CONR_{n1}R_{n2}$, - $C(=N)NR_{n1}R_{n2}$, - CO_2R_o , - $SO_2NR_{n1}R_{n2}$, - SO_3R_o , - SO_2R_o , - $PO(OR_o)_2$, - $NR_{n1}CSNR_{n2}R_{n3}$, - $NR_{n1}C(=N)NR_{n2}R_{n3}$, - $NR_{n1}CONR_{n2}R_{n3}$, - $NR_{n1}COR_c$ and - $NR_{n1}S(=O)_2R_s$;

$R_{n1}, R_{n2}, R_{n3}, R_o$ and R_s are each separately selected from the group consisting of a hydrogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl and may constitute parts of an aliphatic or aromatic heterocycle;

R_c is selected from the group consisting of a hydrogen atom, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, perhalogenated alkyl, cyloalkyl, substituted cycloalkyl, aryl, substituted aryl, benzyl, heteroaryl, substituted heteroaryl, cyano and may constitute parts of an aliphatic or aromatic homo- or heterocycle;

A is selected from the group consisting of an oxygen atom, a sulfur atom, SO , SO_2 , $C(CH_3)_2$ and $C(CF_3)_2$;

E and E' are separately selected from the group consisting of an oxygen atom, a sulfur atom and NR_{n1} ;

G is selected from the group consisting of an oxygen atom, a sulfur atom, and $NR_{n1}R_{n2}$, wherein if G is selected from $NR_{n1}R_{n2}$, G and R_c , as well as G and R_d , may constitute parts of a heterocycle; and

T is selected from the group consisting of an oxygen atom and NR_{n1} .

2. The compound of Claim 1, wherein Y is selected from the group consisting of saturated C_1 - C_{20} alkyl, unsaturated C_1 - C_{20} alkenyl, unsaturated C_1 - C_{20} alkynyl, substituted saturated C_1 - C_{20} alkyl, substituted unsaturated C_1 - C_{20} alkenyl, substituted unsaturated C_1 - C_{20} alkynyl, C_1 - C_{20} cycloalkyl, C_1 - C_{20} cycloalkenyl, substituted saturated C_1 - C_{20} cycloalkyl, substituted unsaturated C_1 - C_{20} cycloalkenyl, aryl, substituted aryl, heteroaryl and substituted heteroaryl.

3. The compound of Claim 2, wherein Y is selected from the group consisting of C₁-C₈ alkyl, C₁-C₈ alkenyl, substituted C₂-C₈ alkyl, substituted C₂-C₈ alkenyl, alkoxyalkyl, aryl, substituted aryl, tertiary and quarternary aminoalkyl, and guanidiniumalkyl groups.

5 4. The compound of Claim 3, wherein Y is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, octyl, and benzyl.

5. The compound of Claim 1, 2, 3 or 4, wherein T is an oxygen atom and G is an a oxygen atom.

10 6. The compound of Claim 1, 2, 3 or 4 wherein m is a positive integer no greater than two.

7. The compound of Claim 1, 2, 3 or 4 wherein m equals one.

8. The compound of Claim 1, 2, 3 or 4 wherein Q is a phenoxide fluorophore.

15 9. The compound of Claim 8, wherein Q is a phenoxide fluorophore selected from the group consisting of a 7-hydroxycoumarin derivative, a resorufin, and a fluorescein.

10. The compound of Claim 1, 2, 3 or 4 wherein R² and R⁴ are hydrogen atoms for all values of p.

11. The compound of Claim 1, 2, 3, 4 or 10, wherein p is equal to one.

20 12. A compound useful as a sensor for cytochrome P450 activity, selected from the group consisting of BOMR, BOMFC, BOMCC, EOMR, EOMCC, MOMCC, MOMFC, MOBFC, OOMR, BOM-DDAO, OOMCC, MOMR, BOM-09B, DBOMF and BOMF, as defined herein.

13. A method of quantifying the activity of a CYP450 enzyme by using the compound of Claim 1, 3, 7, 9, 11 or 12.

25 14. A method for screening a candidate compound for activity as a substrate of at least one CYP450 enzyme, comprising the steps of:

contacting a CYP450 enzyme with the candidate compound and a reagent compound according to Claim 1, 3, 7, 9, 11 or 12; and

30 detecting an optical signal, if any, resulting from interaction of the reagent compound with the CYP450 enzyme.

15. A method for screening a candidate compound for CYP450 inhibitory activity, comprising the steps of:

contacting the candidate compound with a CYP450 enzyme;
contacting the CYP450 enzyme with a reagent compound according to
Claim 1, 3, 7, 9, 11 or 12; and

detecting an optical signal, if any, resulting from interaction of the reagent
compound with the CYP450 enzyme.

16. A method for screening a candidate compound, comprising:
a step for assaying the efficacy of the candidate compound,
a step for assaying the toxicity of the candidate compound, and
a step for assaying the activity of the candidate compound as a substrate or
as an inhibitor of at least one CYP450 enzyme using a reagent compound according
to Claim 1, 3, 7, 9, 11 or 12.

17. The method of Claim 15 or 16 wherein the reagent compound interacts with
the CYP450 enzyme as a competitive or non-competitive inhibitor.

18. The method of Claim 14, 15 or 16 wherein the method comprises a high
throughput screening of a library of fluorogenic candidate compounds.

19. The method of Claim 13, 14 or 15 wherein the CYP450 enzyme is a human
CYP450 enzyme.

20. The method of Claim 14, 15 or 16 wherein the candidate compound is a
member of a library of drug derivatives.

21. The method of Claim 13, 14, 15 or 16 wherein the CYP450 enzyme is
selected from the group consisting of CYP 3A4, CYP 2D6, CYP 2C9, CYP 2C19, CYP
2C8, CYP 2A1, CYP 2B6, CYP 2E1, and CYP 1A2.

22. A candidate drug selected according to the method of Claim 14, 15, 16, 18,
19, 20 or 21.

23. A compound useful as a sensor for cytochrome P450 enzyme activity,
substantially as described herein.

24. A method of quantifying the activity of a CYP450 enzyme, screening a
candidate compound for activity as a substrate of at least one CYP450 enzyme, or
screening a candidate compound for activity as an inhibitor of at least one CYP450
enzyme, by using the compound of Claim 23.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/29439

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D311/16 C07D493/10 C07D311/82 C07D265/38 C12Q1/26

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>MACE ET AL: "Development of CYP450-expressing human bronchial epithelial cell lines for in vitro pharmacotoxicologic applications" IN VITRO TOXICOLOGY, US, MARY ANN LIEBERT, NEW YORK, NY, vol. 10, no. 1, 21 March 1997 (1997-03-21), pages 85-92-92, XP002115754 ISSN: 0888-319X page 85 -page 89</p> <p style="text-align: center;">— — — — — — / —</p>	1,2, 12-24



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

11 April 2000

Date of mailing of the international search report

03/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/29439

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	R.T. MAYER: "7-ALKOXYQUINOLINES: NEW FLUORESCENT SUBSTRATES FOR CYP450" BIOCHEMICAL PHARMACOLOGY, vol. 40, no. 5, 1 September 1990 (1990-09-01), pages 1645-55, XP000901164 OXFORD page 1645 -page 1654	1-4, 12-24
X	IAN N.H. WHITE: "A CONTINUOUS FLUOROMETRIC ASSAY FOR CYP-450" ANALYTICAL BIOCHEMISTRY, vol. 172, no. 2, 1988, pages 304-310, XP000901156 page 304 -page 310	1-4, 12-24
X	JEROEN BUTERS: "A HIGHLY SENSITIVE TOOL FOR THE ASSAY OF CYP450 ENZYME" BIOCHEMICAL PHARMACOLOGY, vol. 46, no. 9, 2 November 1993 (1993-11-02), pages 1577-84, XP000901155 OXFORD page 1577 -page 1582	1-4, 12-24
X	YOSHIAKI OMATA: "CONFORMATIONAL CHANGE OF CYP-450" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 870, no. 3, 1986, pages 392-400, XP000901231 page 392 -page 393	1-4, 12-24
X	MARY HAASCH: "USE OF 7-ALKOXYPHENOXAZONES, 7-ALKOXYCOUMARINS A. 7-ALKOXYQUINOLINES AS FLUORESCENT SUBSTRATES" BIOCHEMICAL PHARMACOLOGY, vol. 47, no. 5, 2 March 1994 (1994-03-02), pages 893-903, XP000901180 OXFORD page 893 -page 899	1-4, 12-24
X	GRANT KRAFFT: "PHOTOACTIVABLE FLUOROPHORES 3." JOURNAL OF THE AMERICAN CHEMICAL SOCIETY., vol. 110, 1988, pages 301-3, XP002135415 AMERICAN CHEMICAL SOCIETY, WASHINGTON, DC., US ISSN: 0002-7863 page 301 -page 303	1,2,8,9, 12
X	GB 2 211 500 A (MEDICAL RESEARCH COUNCIL) 5 July 1989 (1989-07-05) page 1 -page 8	1-4,12
	-/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/29439

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 87 03541 A (POLAROID) 24 September 1987 (1987-09-24) page 31 -page 32; claims	1,2,12
X	EP 0 363 793 A (BASF) 18 April 1990 (1990-04-18) page 1 -page 13	1,2,8,9
X	EP 0 110 682 A (SYVA) 13 June 1984 (1984-06-13) the whole document	1-4,12
X	WO 97 39064 A (MOLECULAR PROBES) 23 October 1997 (1997-10-23) page 0; claim 3	1,2,8,9, 12
X	EP 0 373 104 A (CIBA-GEIGY) 13 June 1990 (1990-06-13) page 1 -page 12	1-4,8,9, 12
P,X	WO 99 58710 A (SMITH-KLINE) 18 November 1999 (1999-11-18) the whole document	1-4,8,9, 12-24
E	WO 00 04008 A (GENTEST CORP.) 27 January 2000 (2000-01-27) the whole document	1-4,8,9, 12-24

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/ 29439

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1

Present claim 1 relates to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds/products/apparatus/methods claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds. The search has been performed among the examples of the description and among the compounds with a similar structure, falling under examples 1 to 8, 20, 21, 30, 31 and tables 1 to 9 and 12 to 14.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/29439

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2211500	A	05-07-1989	NONE	
WO 8703541	A	18-06-1987	US 5196297 A	23-03-1993
			CA 1296010 A	18-02-1992
			CA 1316740 A	27-04-1993
			EP 0250558 A	07-01-1988
			JP 2628607 B	09-07-1997
			JP 7010869 A	13-01-1995
			JP 6098835 B	07-12-1994
			JP 63501941 T	04-08-1988
			US 4970309 A	13-11-1990
			US 5028725 A	02-07-1991
			US 5220036 A	15-06-1993
EP 363793	A	18-04-1990	DE 3834861 A	19-04-1990
			AT 88468 T	15-05-1993
			CA 1336001 A	20-06-1995
			JP 2157273 A	18-06-1990
			JP 2988944 B	13-12-1999
			US 5100914 A	31-03-1992
EP 110682	A	13-06-1984	US 4857455 A	15-08-1989
			AT 66492 T	15-09-1991
			CA 1338608 A	24-09-1996
			DE 3382380 A	26-09-1991
			JP 1950423 C	10-07-1995
			JP 6055158 B	27-07-1994
			JP 59106300 A	19-06-1984
WO 9739064	A	23-10-1997	AU 2801297 A	07-11-1997
			CA 2222275 A	23-10-1997
			EP 0853647 A	22-07-1998
			JP 11508277 T	21-07-1999
EP 373104	A	13-06-1990	CA 2002978 A	17-05-1990
			DE 58908041 D	18-08-1994
			JP 2191673 A	27-07-1990
			JP 2776589 B	16-07-1998
			US 5055595 A	08-10-1991
WO 9958710	A	18-11-1999	NONE	
WO 0004008	A	27-01-2000	NONE	

THIS PAGE BLANK (USPTO)